## PLSC 731: Paper Review, Association Mapping

## Tommasini et al(2007) Association mapping of Stagnospora nodorum blotch resistance in modern European winter wheat varieties. Theor Appl Genet 115:697

- 1. What are the drawbacks to the traditional biparental QTL mapping approach? How can association mapping (AM) address some of these concerns? (697-698)
- 2. What are the differences in mapping resolution between biparental and AM populations? What effect does population size have upon resolution? (698)
- 3. Why is linkage disequilibrium (LD) shorter in an AM population than a biparental population of equivalent size? (698)
- 4. What pleiotropic effects are associated with SNG resistance? (698)
- 5. What have previous SNG studies revealed about the genetic factors controlling resistance? (698)
- 6. Describe the genetic material used in this study? Is it appropriate? (699)
- 7. Are you satisfied with the SNG phenotyping description? (699)
- 8. What type of QTL analysis was performed? (699)
- 9. What molecular markers were used in this study? (699)
- 10. What is the basis of defining subpopulation structure by the STRUCTURE software? (701)
- 11. How was the marker-trait association test significance level set? (701)
- 12. Describe the QTL mapping results. (702)
- 13. Describe the population structure. What criterion was used to assign cultivar to a subpopulation? What geographic feature was observed for the subpopulation? (702)
- 14. Describe the results of the AM analysis. How does the result correlate with the QTL analysis? (703)
- 15. Compare LD in the AM and RIL populations. (704)
- 16. How does the diversity within the AM population compare to the elite European winter wheat germplasm? (704)
- 17. Relative to the marker number and distribution, how might this experiment be improved? Why might a SNG resistant cultivar not have a sun2-3B Arina allele? (705)
- 18. What are the limitations of RIL mapping relative to AM? (706)
- 19. How does the LD testing suggest the AM population was appropriate? (706)
- 20. What is required to better estimate LD in wheat? (706)