

Soybean

Growth and Management

QUICK GUIDE

Reviewed by

Greg Endres, Area Agronomist
Hans Kandel, Agronomist
NDSU Extension Service

Growth, development and yield of soybeans are a result of a variety's genetic potential interacting with environmental and farming practices. Correct production decisions using plant growth staging and timing are important for successful soybean production. Minimizing environmental stress will optimize seed yield.

Farmers who understand how a soybean plant grows and develops can establish their field practices to maximize the genetic potential of the varieties grown. Management practices that may influence crop growth include seedbed preparation, variety selection, planting rate, planting depth, row width, pest management (diseases, insects and weeds), fertilization and harvesting.




Figure 1.
Soybean plant.

NDSU EXTENSION SERVICE

North Dakota State University
Fargo, North Dakota

MAY 2015

Soybeans are classified as indeterminate, semideterminate or determinate in growth in the U.S. Many southern varieties are determinate in growth and cease vegetative growth when the main stem terminates in a cluster of mature pods.

Most northern varieties are indeterminate in growth habit. Indeterminate varieties develop leaves and flowers simultaneously throughout a portion of their reproductive period, with one to three pods at the terminal apex. With soybean development being driven by temperature and photoperiod (interval in a 24-hour period during which a plant or animal is exposed to light), northern varieties have vegetative growth limited by the season length.

Soybean maturity groups are based on adaptation within certain latitudes. These maturity belts run east to west in the U.S., with only about 100 to 150 miles from the north to the south of each belt. Maturity groups range from 000 in the extreme northern U.S. to VIII in the southern Gulf Coast states and most of Florida.

Shortening day length (increasing night length) and warm temperatures control soybean flowering. Soybeans must reach at least the first trifoliolate in growth before they can be induced to flower. Soybeans in the northern U.S. have long minimum day-length requirements for the onset of flowering (often greater than 14 hours of daylight). However, even within a variety, variations in time of flowering may occur from year to year with the same day length closely associated with temperature conditions.

Planting a specific variety farther north than its adapted maturity range will extend the period of vegetative growth, and delay flowering and maturity due to the extended summer day length and cooler temperatures. Likewise, planting a variety farther south than its adapted range will shorten the vegetative growth period, cause earlier flowering and result in an earlier maturity due to shorter summer day length and warmer temperatures.

Vegetative Stages

- VE (emergence)
- VC (cotyledon stage)
- V1 (first trifoliolate)
- V2 (second trifoliolate)
- V3 (third trifoliolate)
- V(n) (nth trifoliolate)
- V6 (flowering will start soon)

Reproductive Stages

- R1 (beginning bloom, first flower)
- R2 (full bloom, flower in top two nodes)
- R3 (beginning pod, 3/16-inch pod in top four nodes)
- R4 (full pod, 3/4-inch pod in top four nodes)
- R5 (1/8-inch seed in top four nodes)
- R6 (full-size seed in top four nodes)
- R7 (beginning maturity, one mature pod)
- R8 (full maturity, 95 percent of pods on the plant are mature)

A fully developed leaf node for the vegetative stages has a leaf above it with unrolled or unfolded leaflets. These unfolded leaflets have their edges no longer touching. Stages are counted from the unifoliolate leaf node upward. All V1 to Vn stages have true leaves that are trifoliolate and produced singularly on different nodes, with these leaves alternating on the stem.

The reproductive stages are divided into four parts: R1 and R2 describe flowering, R3 and R4 describe pod development, R5 and R6 describe seed development and R7 and R8 describe plant maturation.

Vegetative Growth Stages

1. Emergence (VE)

Soybean seed begins germination when the water absorbed is equal to about **50 percent of the seed's weight**. The radical, or primary, root is first to emerge from the seed. Shortly afterward, the hypocotyl (stem) emerges and begins growing toward the soil surface, pulling the cotyledons (seed leaves) with it. This hook-shaped hypocotyl straightens out once emerged and as the cotyledons unfold.

Emergence (VE) normally takes five to 10 days, depending on temperature, moisture conditions, variety and planting depth. During this time, lateral roots also are beginning to grow from the primary root. Root hairs can be visible within five days of planting and provide the key nutrient and water-absorbing functions of the plant in this early stage.

The taproot also will continue growing and branching so that lateral roots can reach the center of a 30-inch row within five to six weeks. Eventually, the soybean root may reach a depth of several feet, with most of the roots in the upper 6 to 12 inches of soil.

Soybeans should be **planted 1 to 1½ inch deep but not deeper than 2 inches**. Because the soybean often must push through crusted soil, deeper planting can limit the viability of seed and final stand number. Rotary hoeing can help break the soil crust, allowing shoots to push through the soil, as well as help in early weed control.

Very small amounts of fertilizer (phosphorus or potassium if needed) in a band to the side and slightly below the seed may help early plant growth, especially if soils are still cool. Do not place fertilizer too near the seed or directly in the furrow because salt injury from the fertilizer can result. **Soybeans are very salt sensitive** (about twice as sensitive as corn).

If fields have **not been in soybeans in the last three years or the field has been flooded**, seed should be inoculated with *Bradyrhizobium japonicum* bacteria to form nodules on the soybean roots that later will provide much of the plant's nitrogen supply. As the hypocotyl arch is exposed to light and straightens to pull the cotyledons out of the ground, growth called epicotyl growth begins with expansion and unfolding of unifoliolate leaves. Auxiliary buds are located where the cotyledons or leaves are attached to the stem.




Figure 2. Germination.

Vegetative Growth Stages

2. Cotyledon Stage (VC)

The VC stage is reached when the unifoliolate leaves are fully expanded. During the VC stage, the **cotyledons supply the nutrient needs** of the young plant (for about seven to 10 days). The cotyledons will lose about 70 percent of their dry weight due to this nutrient reallocation. If one cotyledon is lost during this time, it has little effect on the plant's growth rate. However, loss of both cotyledons at or soon after VE will reduce yields 8 to 9 percent.

Later, after V1, photosynthesis in the developing leaves allows the plant to sustain itself. New **V stages will begin appearing about every three to five days** through V5 and then every two to three days from V5 to shortly after R5, when the node number usually reaches a maximum.

3. First trifoliolate (V1)

This stage is achieved when the first trifoliolate is fully emerged and opened. The V stages after VC are defined and numbered by the upper, fully developed leaf node on the main stem above the unifoliolate leaves (the V1 to Vn stages are numbered by fully developed trifoliolates). Trifoliolate leaves on branches are not counted when determining V stages; only the trifoliolates off the main stem are used in the count.



Figure 3. V2 soybean.

4. Second node (V2)

Plants are 6 to 8 inches tall and have two nodes above the unifoliolate node, with trifoliolate leaves having unfolded leaflets. **Active nitrogen fixation** from the bacteria is just beginning to occur. Most of these root nodules are within 10 inches of the soil surface, with millions of bacteria in each nodule.

Nodules that are pink or red inside are active in nitrogen fixation. White, brown or green nodules are not efficiently fixing nitrogen and are probably parasitic on the plant. The plant will use bacteria-produced nitrogen and the applied or intrinsic soil nitrogen. Too much nitrogen fertilizer will cause the plant to use this supply rather than develop nodules and fix nitrogen.

The top 6 inches of soil have soybean lateral roots developing rapidly. Any **cultivation for weed control should be no deeper than absolutely necessary** to minimize root pruning.

Vegetative Growth Stages

5. Third to Fifth nodes (V3-V5)

Soybean plants are about 7 to 9 inches tall, with three nodes above the unifoliolate node at V3, and will be about 10 to 12 inches tall with five nodes of fully expanded leaflets at V5. The number of branches seen on the plant may increase at this point in wider row spacings and under lower plant densities, depending upon variety grown. Up to six branches are normally developed under field conditions in branched (rather than upright) varieties, with the largest branch being the main stem.

Normally at V5, the plant's axillary buds in the top stem will develop into flower clusters (racemes). **V5 is about one week from R1, or first flower.** At V5, the total number of nodes that the plant can produce is established. With indeterminate soybeans, this total is higher than the number of nodes that actually will develop fully. However, these extra axillary buds allow the plant to recuperate from any hail or wind that may damage some of the growing points.

Although the stem apex (main growing point) is dominant, damage to this growing point will allow axillary buds lower on the plant to branch and grow profusely. Thus, soybeans are capable of producing new branches and leaves after hail destroys almost all of the above-ground foliage as long as at least one axillary bud remains intact. **However, if the plant is broken off below the cotyledonary node, the plant is killed** because no axillary buds are below this node.



Figure 4. V5 soybean.

6. Sixth node (V6)

Soybeans are often 12 to 14 inches tall at this stage, with six nodes containing trifoliolate leaves with fully unfolded and expanded leaflets. The unifoliolate and cotyledons may have senesced from the plant. New stages are unfolding quickly, every two to three days. Lateral roots have crossed over the row underground in any rows 30 inches or less. The plant still is capable of recovering from damage; a **50 percent leaf loss at this stage will affect yield only about 3 percent.**



Figure 5. V6 soybean.

Reproductive Stages




Figure 6. R1 soybean.



Figure 7. R2 soybean.



Figure 8. R3 soybean.

1. Beginning Bloom (R1)

At least one flower is on the plant at any node on the main stem. Soybean **flowering always initiates on the third to sixth node** on the main stem, depending on vegetative stage when flowering begins. This flower initiation will progress up and down the plant. Branches eventually also flower. Within each raceme, the flowering will occur from the base to the tip, so basal pods are always more mature.

Once again, dominance of the primary racemes is seen over secondary racemes on the plant; however, secondary racemes can develop just to the side of primary racemes on the same axil. Vertical roots are rapidly growing and will continue until R4 to R5, as are secondary roots and root hairs nearer the soil surface.

2. Full Bloom (R2)

An open flower is seen at one of the two top nodes of the main stem. At least one of these two upper nodes shows a fully developed leaf. At this stage, **the soybean has accumulated about 25 percent of its total dry weight and nutrients** and has obtained about **50 percent of its mature height**. About 50 percent of the total mature node number has been established. Very rapid nitrogen (N), phosphorus (P), potassium (K) and dry-matter accumulation is occurring and will continue through R6.

The appearance of new flowers on the plant begins to slow between R2.5 and R3 and will be complete by R5. Major lateral roots have turned downward in the soil and nitrogen fixation by root nodules is increasing rapidly. **Defoliation of the plant of 50 percent at this stage will reduce yield by 6 percent.**

3. Beginning Pod (R3)

A pod on the upper four nodes is 3/16 inch long. **Temperature or drought stress at this time can affect yield through total pod number, bean number per pod or seed size.** Partial compensation with only temporary stress can occur in soybeans, but as the plant matures from **R1 to R5.5, this ability to compensate will decrease.**

Very favorable conditions will result in greater pod numbers per plant at this time. Because **60 to 75 percent of most flowers typically abort** on soybeans, any stress that increases this abortion will influence yield greatly. Half of most flowers are lost before pods begin developing and loss of the other half is due to pod abortion. However, the long flowering period of soybeans is one reason the plants can compensate so well.

Reproductive Stages

4. Full Pod (R4)

This stage shows rapid pod growth and the beginning of seed development at the beginning of the full-pod stage. The dry weight of pods is increased greatly from R4 to R5. The plant has a pod at this stage that is at least 3/4 inch long on at least one of the four upper nodes of the main stem. **This stage is the most crucial period for seed yield.** Any stress from R4 to R6 causes more yield reduction than at any other time.

Late pod formation at R4.5 to early seed fill at R5.5 is most critical. Yield reduction at this time is mainly from fewer pods. This is a critical period to consider irrigation, if needed, to reduce yield losses. The last flowering will occur at the main stem tip (through R5).



Figure 9. R4 soybean.

5. Beginning Seed (R5)

Seed filling during this stage requires much water and nutrients from the plant. Redistribution of nutrients in the plant occurs with the soybean providing about a half of needed N, P and K from the plant's vegetative parts and about a half from N fixation and nutrient uptake by the roots.

Leaf loss of 100 percent at this stage will reduce yields by 80 percent; the plant is less able to compensate from stress and vegetative damage. Stress can lower yields by reducing pod numbers and the number of beans per pod, and to a lesser extent, by reducing seed size.

This stage has seed at least 1/8 inch long in one of the pods on one of the four upper nodes of the main stem. **About halfway through this stage, the plant attains its maximum height, node number and leaf area.** Nitrogen fixation peaks and begins to drop, and the seeds continue a steady period of dry weight accumulation. Toward the end of this stage, the nutrient accumulation in the leaves peaks and then begins the process of redistributing to the seed.

Seed accumulation will continue until shortly after R6.5, with about 80 percent of total seed dry weight being accomplished.



Figure 10. R5 soybean.

6. Full Seed (R6)

This stage also is known as the "green bean" stage or beginning full-seed stage, and total pod weight will peak during this stage. The growth rate of the beans is rapid but will slow by R6.5 and peak at R7.

This stage initiates with a pod containing a green seed that fills the pod cavity on at least one of the four top nodes of the main stem. Rapid leaf yellowing will begin right after this stage until R8, or all leaves have fallen. Within this stage, three to six trifoliolate leaves may fall from the lowest nodes on the plant prior to leaf yellowing. **Root growth is complete at about R6.5.**



Figure 11. R6 soybean.

Reproductive Stages




Figure 12. R7 soybean.




Figure 13. R8 soybean.

7. Beginning Maturity (R7)

This stage begins with one normal pod on the main stem, which obtains the mature color (brown or tan). **Dry matter begins to peak in individual seeds.** This is visible when all the green is lost from the seeds and pods (they appear yellow).

Seeds contain about 35 percent moisture at physiological maturity. Stress at this stage or later has almost no effect on yield unless pods are dropped to the ground or seeds are shattered from the pods. Also, any lodged plants may reduce actual yield (due to reduced light interception) and harvested yield (harvest losses). At this stage, the crop is safe from a killing frost.

8. Full Maturity (R8)

On the soybean plant, 95 percent of the pods have reached their mature color and **only five to 10 days of good drying weather after this stage will be required to have the soybeans at less than 15 percent moisture**, or harvest moisture. Soybeans will lose moisture rapidly with warm and dry weather at this point but should be harvested soon to prevent losses.

Long-term storage of soybeans should be at 13 percent or less moisture. When harvesting, leave short stubble to ensure a full harvest. **A 3½-inch stubble still leaves 5 percent or more of the crop and a 6½-inch stubble, 12 percent or more.**

This publication was authored by D.R. Berglund, professor emeritus and former Extension agronomist; D.A. McWilliams, former Extension agronomist; and G.J. Endres, Extension area agronomist, NDSU, 1999.

Photography for this publication was provided through the Iowa State University Photo Service, J.C. Herman, editor. The stage designation format was derived from Iowa State Extension special report No. 53, "How a Soybean Plant Develops," prepared by S.W. Ritchie, J.J. Hanway, H.E. Thompson and G.O. Benson. Also, qualification on production management suggestions were summarized from lectures by J.R. George through Iowa State University. Special thanks to T. Helms, professor, NDSU, for suggestions and editorial review.

The NDSU Extension Service does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names. NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

For more information on this and other topics, see www.ag.ndsu.edu

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. North Dakota State University does not discriminate on the basis of age, color, disability, gender expression/identity, genetic information, marital status, national origin, public assistance status, race, religion, sex, sexual orientation, or status as a U.S. veteran. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708. This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.

3M-6-99, 1M-5-01, 1M-8-04, 1M-5-15