
A1145



R-5.1



R-7



R-8

R-9



R-6

North Dakota State University Fargo, North Dakota

Reviewed July 2019





Description of sunflower growth stages

The total time required for development of a sunflower plant and the time between the various stages of development depends on the genetic background of the plant and the growing environment. When determining the growth stage of a sunflower field, the average development of a large number of plants should be considered. This staging method also can be used for individual plants. The same system can be used for classifying a single head or branched sunflower. In the case of branched sunflower, make determinations using only the main branch or head. In stages R7 through R9, use healthy, disease-free heads to determine plant development if possible because some diseases can cause head discoloration.

	Stage	Description
Vegetative Emergence	VE	Seedling has emerged and the first leaf beyond the cotyledons is less than 4 cm long.
Vegetative Stages	V (number) (i.e.) V1 V2 V3 etc.,	These are determined by counting the number of true leaves at least 4 cm in length beginning as V1, V2, V3, V4, etc. If senescence of the lower leaves has occurred, count leaf scars (excluding those where the cotyledons were attached) to determine proper stage.
Reproductive Stages	R1	The terminal bud forms a miniature floral head rather than a cluster of leaves. When viewed from directly above, the immature bracts form a many-pointed starlike appearance.
	R2	The immature bud elongates 0.5 to 2 cm above the nearest leaf attached to the stem. Disregard leaves attached directly to the back of the bud.
	R3	The immature bud elongates more than 2 cm above the nearest leaf.
	R4	The inflorescence begins to open. When viewed from directly above, immature ray flowers are visible.
	R5 (decimal) (i.e.) R5.1 R5.2 R5.3 etc.	This stage is the beginning of flowering. The stage can be divided into substages dependent upon the percent of the head area (disk flowers) that has completed or is in flowering. Ex. R5.3 (30%), R5.8 (80%) etc.
	R6	Flowering is complete and the ray flowers are wilting.
	R7	The back of the head has started to turn pale yellow.
	R8	The back of the head is yellow but the bracts remain green.
	R9	The bracts become yellow and brown. This stage is regarded as physiological maturity.

From Schneiter, A.A., and J.F. Miller. 1981. Description of Sunflower Growth Stages. Crop Sci. 21:901-903.

NDSU Extension does not endorse commercial products or companies even though reference may be made to tradenames, trademarks or service names. NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

For more information on this and other topics, see www.ndsu.edu/extension

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost for Title IX/ADA Coordinator, Old Main 201, NDSU Main Campus, 701-231-7708, ndsu.eoaa@ndsu.edu. This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881. 2M-2-98, 1.5M-4-05, 300-5-13