

Older Adults and Food Safety: **To Market, To Market**

Preventing Foodborne Illness

Prevention of foodborne illness starts with your trip to the supermarket. Pick up your packaged and canned foods first.

Don't buy food in cans that are bulging or dented or jars that are cracked or have loose or bulging lids. Look for expiration dates on the labels and never buy outdated food.

Likewise, check the "use by" or "sell by" dates on dairy products, such as cottage cheese, cream cheese, yogurt and sour cream, and pick the ones that will stay fresh longest in your refrigerator.

Choose eggs that are refrigerated in the store. Before putting them in your cart, open the carton to make sure none are cracked or leaking.

Save frozen foods and perishables such as meat, poultry or fish until last. Always put these products in separate plastic bags so that drippings don't contaminate other foods in your shopping cart.

Check for cleanliness at the meat or fish counter and the salad bar. For instance, cooked shrimp lying on the same bed of ice as raw fish could be contaminated. When buying from a salad bar, avoid fruits and vegetables that look brownish, slimy or dried out. These are signs that the product has been held at an improper temperature.

When shopping for shellfish, buy from markets that get their supplies from state-approved sources; stay clear of vendors who sell shellfish from roadside stands or the back of a truck. And if you're planning to harvest your own shellfish, heed posted warnings about the safety of the water.

Take an ice chest along to keep frozen and perishable foods cold if getting your groceries home will take more than two hours.



NDSU EXTENSION

North Dakota State University Fargo, North Dakota

Safe Handling of Fruits and Vegetables

- Wash hands with warm water and soap for at least 20 seconds before and after handling food, especially fresh, whole fruits and vegetables, and raw meat, poultry and fish. Clean under fingernails, too.
- Rinse raw products in warm water. Don't use soap or other detergents. If necessary and appropriate, use a small scrub brush to remove surface dirt.
- Use smooth, durable and nonabsorbent cutting boards that can be cleaned and sanitized easily.
- Wash cutting boards with hot water, soap and a scrub brush to remove food particles. Then sanitize the boards by putting them through the automatic dishwasher or rinsing them in a solution of 1 teaspoon of chlorine bleach to 1 quart of water. Always wash boards and knives after cutting raw meat, poultry or seafood and before cutting another food to prevent cross-contamination.
- Store cut, peeled and brokenapart fruits and vegetables (such as melon balls) at or below 41 F in the refrigerator.

Julie Garden-Robinson, Ph.D., R.D., L.R.D. Food and Nutrition Specialist

Source: Adapted with permission from FDA/Center for Food Safety and Applied Nutrition, May 1999; www.cfsan.fda.gov/~dms/seniorsd.html Developed in cooperation with AARP This material is based upon work supported by the Cooperative State Research, Education and Extension Service, U.S. Department of Agriculture, under Agreement No. 2002-51110-01512.

Apply the Heat and Fight BAC!

Cooking food to the proper temperature kills harmful bacteria. This includes raw meat, poultry, fish and eggs, as well as foods that are thoroughly cooked upon purchase. Precooked foods may become contaminated during storage or handling.

Reheat ready-to-eat foods, such as hot dogs, luncheon meats, cold cuts, fermented and dry sausage, and other deli-style meat and poultry products, until they are steaming hot. If you cannot reheat these foods, do not eat them.

Cooking food to the proper temperature kills harmful bacteria. So Fight BAC![™] by thoroughly cooking your food as follows:

Raw Food	Internal Temperature
Ground Products	
Hamburger	160 F
Beef, veal, lamb, pork	160 F
Chicken, turkey	165 F
Beef, Veal, Lamb	
Roasts and steaks	
- medium-rare	145 F*
- medium	160 F
- well-done	170 F
Pork	
Chops, roasts, ribs	
- medium-rare	145 F*
- medium	160 F
- well-done	170 F
Ham, fresh	145 F*
Sausage, fresh	160 F
Poultry	
Chicken, whole and pieces	165 F
Duck	165 F
Turkey (unstuffed)	165 F
- Whole	165 F
- Breast	165 F
- Dark meat	165 F
- Stuffing (cooked separate	<i>ly)</i> 165 F
Eggs	
	hite are firm
Casseroles	160 F
Sauces, custards	160 F

*Allow three-minute rest time.

This chart has been adapted for home use and is consistent with consumer guidelines from the U.S. Department of Agriculture and U.S. Food and Drug Administration (FDA).

For more information about food safety, visit the NDSU Extension website: www.ag.ndsu.edu/food

County commissions, North Dakota State University and U.S. Department of Agriculture cooperating. NDSU does not discriminate in its programs and activities on the basis of age, color, gender expression/identity, genetic information, marital status, national origin, participation in lawful off-campus activity, physical or mental disability, pregnancy, public assistance status, race, religion, sex, sexual orientation, spousal relationship to current employee, or veteran status, as applicable. Direct inquiries to Vice Provost for Title IX/ADA Coordinator, Old Main 201, NDSU Main Campus, 701-231-7708, ndsu.eoaa@ndsu.edu. This publication will be made available in alternative formats for people with disabilities upon request, 701-231-7881.