

# Mid-Miocene cooling and the extinction of tundra in continental Antarctica

Adam R. Lewis<sup>a,b,c,d</sup>, David R. Marchant<sup>a</sup>, Allan C. Ashworth<sup>c</sup>, Lars Hedenäs<sup>e</sup>, Sidney R. Hemming<sup>f</sup>, Jesse V. Johnson<sup>g</sup>, Melanie J. Leng<sup>h</sup>, Malka L. Machlus<sup>f</sup>, Angela E. Newton<sup>i</sup>, J. Ian Raine<sup>j</sup>, Jane K. Willenbring<sup>k</sup>, Mark Williams<sup>l</sup>, and Alexander P. Wolfe<sup>m</sup>

<sup>a</sup>Department of Earth Sciences, Boston University, Boston, MA 02215; <sup>b</sup>Byrd Polar Research Center, Ohio State University, Columbus, OH 43210; <sup>c</sup>Department of Geosciences, North Dakota State University, Fargo, ND 58105; <sup>d</sup>Department of Cryptogamic Botany, Swedish Museum of Natural History, Box 50007, SE-104 05 Stockholm, Sweden; <sup>e</sup>Lamont–Doherty Earth Observatory, Columbia University, Palisades, NY 10964; <sup>f</sup>Department of Computer Science, University of Montana, Missoula, MT 59812; <sup>g</sup>Natural Environment Research Council Isotope Geosciences Laboratory, Keyworth, Nottingham NG12 5GG, United Kingdom; <sup>h</sup>Department of Botany, Natural History Museum, London SW7 5BD, United Kingdom; <sup>i</sup>GNS Science, Lower Hutt 5040, New Zealand; <sup>j</sup>Institut für Mineralogie, Leibniz Universität, D-30167 Hannover, Germany; <sup>k</sup>Department of Geology, University of Leicester, Leicester LE1 7RH, United Kingdom; and <sup>m</sup>Department of Earth and Atmospheric Sciences, University of Alberta, Edmonton, AB, Canada T6G 2E3

Edited by James P. Kennett, University of California, Santa Barbara, CA, and approved June 3, 2008 (received for review March 12, 2008)

**A major obstacle in understanding the evolution of Cenozoic climate has been the lack of well dated terrestrial evidence from high-latitude, glaciated regions. Here, we report the discovery of exceptionally well preserved fossils of lacustrine and terrestrial organisms from the McMurdo Dry Valleys sector of the Transantarctic Mountains for which we have established a precise radiometric chronology. The fossils, which include diatoms, palynomorphs, mosses, ostracodes, and insects, represent the last vestige of a tundra community that inhabited the mountains before stepped cooling that first brought a full polar climate to Antarctica. Paleocological analyses, <sup>40</sup>Ar/<sup>39</sup>Ar analyses of associated ash fall, and climate inferences from glaciological modeling together suggest that mean summer temperatures in the region cooled by at least 8°C between 14.07 ± 0.05 Ma and 13.85 ± 0.03 Ma. These results provide novel constraints for the timing and amplitude of middle-Miocene cooling in Antarctica and reveal the ecological legacy of this global climate transition.**

climate change | tundra biota | Dry Valleys | diatoms | ostracodes

The Earth's climate has been cooling for the last 50 million years in a trend punctuated by three steps. Higher rates of cooling occurred during the late Eocene–early Oligocene (c. 34 Ma), the middle Miocene (c. 14 Ma), and the late Pliocene (c. 3 Ma) (1). The step at 34 Ma is associated with the onset of continental glaciation in Antarctica, and that centered on 3 Ma registers the expansion of Northern Hemisphere ice sheets. The mid-Miocene step, often referred to as the middle-Miocene climatic transition (MMCT), is the least understood, with significant questions surrounding its timing, style, and fundamental cause. Marine isotopic studies imply significant ice volume expansion in Antarctica and coeval cooling of the Southern Ocean (2). Because of the paucity of dated terrestrial archives from the continent, little evidence for climate change has been available to constrain directly the nature and timing of the MMCT. This has resulted in a number of unanswered questions. To what extent did the continent cool during the MMCT, and was this cooling gradual or abrupt? What effect did the MMCT have on the thermal regime of Antarctic glaciers? And from a biological perspective, what was the ecological legacy of the MMCT for organisms whose ancestors had survived the cooling step at 34 Ma? Did the extinction of a shrub tundra occur during the middle Miocene, or was it delayed until the Pliocene (3, 4)?

The fossiliferous glaciogenic sediments reported here are from the western Olympus Range in the McMurdo Dry Valleys sector of the Transantarctic Mountains. The fossils and their glacial geologic context permit analyses of the MMCT from three broad perspectives: paleocological, glaciological, and geomorphological. The *in situ* fossil-bearing strata occur within several north-facing valleys that open into McKelvey Valley, one of the major

ice-free troughs of the McMurdo Dry Valleys (Fig. 1). Prior work has shown that surficial deposits in the Range include a basal group of unconsolidated tills deposited from wet-based alpine glaciers and an overlying group of stacked sublimation tills from cold-based alpine glaciers; on the basis of dated ashfall covering the oldest (stratigraphically lowest) sublimation till, the transition from wet- to cold-based glaciation occurred before 13.85 ± 0.03 Ma (5). Results from our recent fieldwork show that lacustrine deposits interfinger with wet-based tills deposited just before the transition, indicating that a series of small lakes developed where recessional moraines impounded meltwater (Fig. 1). It is within these lacustrine sediments that we find well preserved freshwater fossils and pristine, bedded ashfall. A date of 14.07 ± 0.05 Ma for an *in situ* ashfall layer, 3 cm thick, within an extensive sequence of locally fossiliferous lake deposits provides a direct, unambiguous age for the last phase of wet-based alpine glaciation and an age for tundra plants and animals that had colonized the locally deglaciated terrain [Fig. 1, [supporting information \(SI\) Table S1](#), and [Fig. S1](#)].

The best-studied fossil assemblage comes from a small (14,000 m<sup>2</sup>) moraine-dammed basin at 1,425 m near Mount Boreas in the western Olympus Range (Fig. 1). Deflation by wind scour has exposed lacustrine beds, which are unconsolidated, planar, and near-horizontal except where locally disturbed by the development of polygonal patterned ground. At the basin center, the stratigraphy records infilling, first by glaciolacustrine silt and sand, later by diatom- and moss-rich muds, and finally by fluvial sands and debris flows (Fig. 2). The diatom stratigraphy suggests that the lake persisted for thousands of years, during which time water levels fell at least once and the floor of the basin became a bryophyte-rich mire. The small alpine lake continued to support obligate freshwater organisms even during the low water phase.

Many of the fossils from the Mount Boreas lacustrine beds are preserved in exquisite detail. The majority represent aquatic taxa that inhabited the lake during an early shallow-water and later deep-water phase. The shallow-water assemblage is represented

Author contributions: A.R.L., D.R.M., A.C.A., L.H., S.R.H., J.V.J., M.J.L., M.L.M., A.E.N., J.I.R., J.K.W., M.W., and A.P.W. performed research; A.R.L. and A.C.A. analyzed data; and A.R.L., D.R.M., A.C.A., and A.P.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.


<sup>d</sup>To whom correspondence should be sent at the present address: Department of Geosciences, North Dakota State University, Fargo, ND 58105. E-mail: adam.r.lewis@ndsu.edu.

This article contains supporting information online at [www.pnas.org/cgi/content/full/0802501105/DCSupplemental](http://www.pnas.org/cgi/content/full/0802501105/DCSupplemental).

© 2008 by The National Academy of Sciences of the USA







**Fig. 4.** Fossil ostracode and diatoms. (A) Soft anatomy preserved between the valves of an ostracode in the superfamily Cypridoidea. (B) Disarticulated colony of the benthic alkaliphilous diatom *Staurosirella pinnata*. (C) *Pinnularia* sp., a benthic periphytic form. (D) *Aulacoseira* sp., a tycho planktonic filamentous diatom. (E) The planktonic form *Asterionella formosa*. (Scale bars: in B–E, 5  $\mu$ m.)

than that of today. We are able to estimate summer warmth from the modern autecologies of several of the recovered taxa. The closest locations to the McMurdo Dry Valleys inhabited by the moss *D. longifolius* today is on James Ross and Vega islands<sup>8</sup> at the northern tip of the Antarctic Peninsula (64°S) where the mean summer temperature (MST) is  $-1.7^{\circ}\text{C}$ .<sup>9</sup> *D. longifolius* is such a widely distributed moss in both the northern and southern hemispheres (18) that we assume it would have colonized farther south in continental Antarctica if there were sufficient warmth and moisture to support its growth. Whereas the moss *D. longifolius* provides an estimate for the minimum thermal environment at the site, the paleoecology of fossil ostracodes and beetle indicate a warmer MST. The nearest lakes with cyprid-

doidean ostracodes today occur in the South Orkney Islands (60°S) where MST is approximately  $1^{\circ}\text{C}$  (19, 20). Likewise, the southernmost records for indigenous beetles are for the island of South Georgia (55°S), where the MST is approximately  $5^{\circ}\text{C}$ .<sup>10</sup> Lathridiid beetle species have been recorded farther south on South Orkney and South Shetland islands (62°S), but they are considered to be recent human introductions (21). Together, the fossil data suggest that MST was  $\approx 5^{\circ}\text{C}$ . In comparison, the MST at the site today is approximately  $-12^{\circ}\text{C}$  (22).

To estimate the amount of cooling that occurred between the wet-based glaciation at  $14.07 \pm 0.05$  Ma and the onset of cold-based glaciation at  $13.85 \pm 0.03$  Ma, we used a steady-state thermomechanically coupled glaciological model with higher-order stress treatments (see *SI Text*). The model provides a test of basal-ice sensitivity for local alpine glaciers to varying atmospheric temperatures. Assumptions include minimal change in the elevation of the western Olympus Range (23) and that the former wet-based alpine glaciers were similar in size and shape to modern cold-based counterparts throughout the McMurdo Dry Valleys region; the latter assumption is borne out by detailed mapping of recessional moraines from wet-based alpine glaciers in the western Olympus Range. Model results show that the maximum atmospheric temperature consistent with cold-based alpine glaciation in the western Olympus Range is  $-3^{\circ}\text{C}$  (Fig. S2). Given that maximum atmospheric temperatures are achieved during summer, the value can be seen as a conservative proxy for MST. If the landscape during the mid-Miocene were at a lower elevation, as some models of landscape evolution imply (24), then an even lower MST would be required to maintain cold-based glaciation. The combined results from paleoecology and glaciological modeling indicate a minimum decrease in MST of  $8^{\circ}\text{C}$  between 14.07 and 13.85 Ma. This interpretation is consistent with the timing of a sea-surface cooling of  $6\text{--}7^{\circ}\text{C}$  between 14.2 and 13.8 Ma inferred from Mg/Ca ratios of Southern Ocean planktonic foraminifera (2).

An ongoing debate centers on when hyperarid, polar-desert conditions became established in Antarctica. At question is the long-term stability of the East Antarctic Ice Sheet and whether global warming during the Pliocene ( $\approx 3$  Ma) (25) resulted in deglaciation of the continental interior and permitted recolonization of tundra organisms as far south as lat.  $85^{\circ}\text{S}$  (4, 26, 27). Our combined paleoglaciological and paleoecological results clarify several of these issues. First, the glacial stratigraphy and geomorphology of the McMurdo Dry Valleys indicate that the transition to cold-based, alpine glacial regimes at 13.85 Ma was never subsequently reversed (5). Second, the exquisite preservation of several fossil groups recovered from Miocene lake sediments (Figs. 3 and 4) indicates minimal postdepositional degradation, which is best reconciled by these assemblages having remained in a perennially dry and frozen state. Third, glass shards from the dated ash deposits show no evidence of either hydration or authigenic clay mineral formation (5, 28), suggesting essentially no chemical weathering. Taken together, these observations constitute robust arguments against the return to warmer and wetter conditions at any time after the MMCT. Accordingly, we surmise that polar desert climate has persisted uninterrupted in the western Olympus Range, and by inference all regions in the Transantarctic Mountains south of  $77^{\circ}\text{S}$ , since at least 13.85 Ma. This climate reconstruction is incompatible with either a marked reduction in the volume of the East Antarctic Ice Sheet inland of the McMurdo Dry Valleys region, or with the recolonization by tundra biota during the middle Pliocene.

<sup>8</sup>British Antarctic Survey Plant Database, [www.antarctica.ac.uk/bas\\_research/data/access/plantdatabase/index.php](http://www.antarctica.ac.uk/bas_research/data/access/plantdatabase/index.php). Last updated 2007; last accessed January 8, 2008.

<sup>9</sup>Antarctic Climate Data: Results from the SCAR Reader Project. [www.antarctica.ac.uk/met/gjma/temps.html](http://www.antarctica.ac.uk/met/gjma/temps.html). Last updated June 3, 2007; last accessed January 8, 2008.

<sup>10</sup>Antarctic CRC and Australian Antarctic Division Climate Data Sets. [www.antarctica.ac.uk/met/READER/temperature.html](http://www.antarctica.ac.uk/met/READER/temperature.html). Last updated February 29, 2000; last accessed January 8, 2008.

The evidence presented here suggests that the climate of East Antarctica cooled abruptly between  $14.07 \pm 0.05$  and  $13.85 \pm 0.03$  Ma. The ultimate cause of the climate change is uncertain. A decline in global  $p\text{CO}_2$ , based on changes in stomatal density on fossil leaves, has been proposed as the primary driver (29), although marine geochemical evidence suggests that  $p\text{CO}_2$  declined several million years earlier than the MMCT (30). Alternative explanations focus on tectonically driven changes to oceanic circulation resulting in a reorganization of the ocean–atmosphere–cryosphere system (2, 31, 32). Regardless of the ultimate cause, this shift was sufficient to initiate the transition from wet- to cold-based alpine glaciation in the central and southern Transantarctic Mountains, and to induce the regional extinction of tundra plants and animals. The only possible survivors from this earlier biota are microscopic, mostly soil-

dwelling organisms (33). The climatic and environmental changes reported here were likely the most significant to have impacted Antarctica since the initial onset of Cenozoic continental glaciation. These results provide challenging but testable targets for climate models, as well as new constraints for Antarctic biogeography.

**ACKNOWLEDGMENTS.** We thank P. Wilby for his work on ostracode taphonomy that recognized the preservation by iron oxide; D. Horne and D. Siveter for advice on ostracode anatomy; B. Vandenheuevel, H. Margerison, S. Burns, E. Klingler, D. Kowalewski, D. Shean, K. Swanger, K. Gorz, and A. Podoll for help in excavating and mapping the lacustrine deposits; R. Thomasson, T. Kummer, and A. Pannem for their help in sample processing; and R. Mink for her illustration of the beetle fossil. This study was supported by National Science Foundation Grants ANT 0440711 (to D.R.M.), ANT 9811877 (to D.R.M.), and ANT-0230696 and ANT 0440761 (to A.C.A.).

- Zachos J, Pagani M, Stone L, Thomas E, Billups K (2001) Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693.
- Shevenell AE, Kennett JP, Lea DW (2004) Middle Miocene Southern Ocean cooling and Antarctic cryosphere expansion. *Science* 305:1766–1770.
- Webb P-N, Harwood DM, McKelvey BC, Mercer JH, Stott LD (1984) Cenozoic marine sedimentation and ice-volume variation on the East Antarctic craton. *Geology* 12:287–291.
- Hill RS, Harwood DM, Webb P-N (1996) *Nothofagus beardmorensis* (Nothofagaceae), a new species based on leaves from the Pliocene Sirius Group, Transantarctic Mountains, Antarctica. *Rev Palaeobot Palynol* 94:11–24.
- Lewis AR, Marchant DR, Ashworth AC, Hemming SR, Machlus ML (2007) Major middle Miocene global change: Evidence from East Antarctica and the Transantarctic Mountains. *Geol Soc Am Bull* 119:1449–1461.
- Spaulding SA, McKnight DM (1999) In *The Diatoms: Applications to the Environmental and Earth Sciences*, eds Stoermer E, Smol J, (Cambridge Univ Press, Cambridge, UK), pp 245–263.
- Kellogg TB, Kellogg DE (2002) *Non-Marine and Littoral Diatoms from Antarctic and Subantarctic Regions: Distribution and Updated Taxonomy*, Diatom Monographs, ed Witkowski A (Gantner, Ruggell, Liechtenstein, Koeltz Scientific, Königstein, Germany), Vol 1.
- Joynt EH, Wolfe AP (2001) Paleoenvironmental inference models from sediment diatom assemblages in Baffin Island lakes (Nunavut, Canada) and reconstruction of summer water temperature. *Can J Fish Aquat Sci* 58:1222–1243.
- Smol JP, et al. (2005) Climate-driven regime shifts in the biological communities of arctic lakes. *Proc Natl Acad Sci USA* 102:4397–4402.
- Whitehead DR, Charles DF, Jackson ST, Smol JP, Engstrom ER (1989) The developmental history of Adirondack (N.Y.) lakes. *J Paleolimnol* 2:185–206.
- Renberg I (1990) A 12,600 year perspective of the acidification of Lilla Oresjon, Southwest Sweden. *Philos Trans R Soc London* 327:357–361.
- Michelutti N, Wolfe AP, Briner JP, Miller GH (2007) Climatically controlled chemical and biological development in Arctic lakes. *J Geophys Res* 112:G03002.
- Engstrom DR, Fritz SC, Almendinger JE, Juggins S (2000) Chemical and biological trends during lake evolution in recently deglaciated terrain. *Nature* 408:161–166.
- Prebble JG, Raine JI, Barrett PJ, Hannah MJ (2006) Vegetation and climate from two Oligocene glacioeustatic sedimentary cycles (31 and 24 Ma) cored by the Cape Roberts Project, Victoria Land Basin, Antarctica. *Palaeogeogr Palaeoecol* 231:41–57.
- Raine JI (1998) Terrestrial palynomorphs from Cape Roberts Project Drillhole CRP-1, Ross Sea, Antarctica. *Terra Antarctica* 5:539–548.
- Morrhead DL, Priscu JC (1998) In *Ecosystem Dynamics in a Polar Desert: The McMurdo Dry Valleys, Antarctica*, AGU Antarctic Research Series, ed Priscu JC (Am Geophys Union, Washington, DC), pp 351–369.
- Stevens MI, Hogg ID (2002) Expanded distributional records of Collembola and Acari in southern Victoria Land, Antarctica. *Pedobiologia* 46:485–495.
- Hedenäs L (1997) The Drepanocladus s. str. species with excurrent costae (Muscic: Amblystegiaceae). *Nova Hedwigia* 64:535–547.
- Pugh PJA, Dartnall HJG, McInnes SJ (2002) The non-marine Crustacea of Antarctica and the Islands of the Southern Ocean: Biodiversity and biogeography. *J Nat Hist* 36:1047–1103.
- Peck LS, Convey P, Barnes KA (2006) Environmental constraints on life histories in Antarctic ecosystems: Tempos, timings and predictability. *Biol Rev* 81:75–109.
- Gressitt JL (1967) In *Entomology of Antarctica*, AGU Antarctic Research Series, ed Gressitt J (Am Geophys Union, Washington, DC), Vol 10, pp 1–33.
- Doran PT, et al. (2002) Valley floor climate observations from the McMurdo Dry Valleys, Antarctica, 1986–2000. *J Geophys Res*, 10.1029/2001JD002045.
- Sugden DE, Denton GH, Marchant DR (1995) Landscape evolution of the McMurdo Dry Valleys, Transantarctic Mountains: Tectonic implications. *J Geophys Res* 100:9949–9967.
- Stern TA, Baxter AK, Barrett PJ (2005) Isostatic rebound due to glacial erosion within the Transantarctic Mountains. *Geology* 33:221–224.
- Naish T, et al. (2007) A record of Antarctic climate and ice sheet history recovered. *Eos* 88:557–558.
- Ashworth AC, Cantrill DJ (2004) Neogene vegetation of the Meyer Desert formation (Sirius Group) Transantarctic Mountains, Antarctica. *Palaeogeogr Palaeoecol* 213:65–82.
- Ashworth AC, Thompson FC (2003) A fly in the biogeographic ointment. *Nature* 423:135.
- Marchant DR, Denton GH, Sugden DE, Swisher CC (1993) Miocene glacial stratigraphy and landscape evolution of the western Asgard Range, Antarctica. *Geogr Ann* 75A:303–330.
- Kürschner WM, Kvaček Z, Dilcher DL (2008) The impact of Miocene atmospheric carbon dioxide fluctuations on climate and the evolution of terrestrial ecosystems. *Proc Natl Acad Sci USA* 105:449–453.
- Pagani M, Freeman KH, Arthur MA (1999) Late Miocene Atmospheric CO<sub>2</sub> Concentrations and the Expansion of C<sub>4</sub> Grasses. *Science* 285:876–879.
- Kennett JP (1975) Cenozoic evolution of Antarctic glaciation, the Circum-Antarctic Ocean and their impact on global paleoceanography. *J Geophys Res* 82:3843–3860.
- Woodruff F, Savin SM (1991) Mid-Miocene isotope stratigraphy in the deep sea: High resolution correlations, paleoclimatic cycles, and sediment preservation. *Paleoceanography* 6:755–806.
- Convey P, Stevens MI (2007) Antarctic biodiversity. *Science* 317:1877–1878.