


# Pfizer COVID-19 Vaccination Schedule for People Who are Moderately or Severely Immunocompromised

		Months 			
Age & Indication	COVID-19 Vax History	0	1	2	3
Pfizer <sup>a</sup> for 6 months - 4 years	Unvaccinated	1st bivalent dose	2nd bivalent dose (3 weeks after 1st)		3rd bivalent dose (>8 weeks after 2nd)
	1 dose of monovalent		1st bivalent dose (3 weeks after last monovalent dose)		2nd bivalent dose (>8 weeks after 1st)
	2 or 3 doses of monovalent			1 bivalent dose (>8 weeks after last monovalent dose)	
	2 or 3 doses of monovalent + 1 dose of bivalent			*OPTIONAL* Additional bivalent dose	
Pfizer <sup>a</sup> for ≥5 years	Unvaccinated	1st bivalent dose	2nd bivalent dose (3 weeks after 1st)	3rd bivalent dose (>4 weeks after 2nd)	
	1 dose of monovalent		1st bivalent dose (3 weeks after last monovalent dose)	2nd bivalent dose (>4 weeks after 1st)	
	2 doses of monovalent			1st bivalent dose (>4 weeks after last monovalent dose)	
	3 doses of monovalent			1st bivalent dose (>8 weeks after last monovalent dose)	
	3 doses of monovalent + 1 dose of bivalent			*OPTIONAL* Additional bivalent dose	

a. For people who are moderately or severely immunocompromised: At the time of initial vaccination, people ages >6 months are recommended to receive 3 bivalent mRNA doses. People ages >6 months who previously received only monovalent doses are recommended to receive 1 or 2 bivalent mRNA vaccine doses depending on age and vaccine product. People who previously received a bivalent mRNA vaccine dose(s) have the option to receive 1+ additional bivalent mRNA doses. Any additional doses must be determined by the clinical judgement of a healthcare provider and personal preference and circumstances. Any further additional doses should be administered at least 2 months after the last COVID-19 vaccine dose.