



**North Dakota  
Water Resources Research Institute**

**North Dakota State University  
University of North Dakota**

**ANNUAL REPORT**

March 1, 2011 to February 29, 2012

**Fiscal Year 2011 Report to the U.S. Geological Survey**

June 2012

# **Annual Report**

## **Fiscal Year 2011 Report to the U.S. Geological Survey**

Award No. 06-HQ-GR-0104  
U.S. Geological Survey

G. Padmanabhan, Director  
North Dakota Water Resources Research Institute  
North Dakota State University  
Fargo, ND 58108-6050

June 2012

## Table of Contents

INTRODUCTION -----	1
Program Management -----	2
State Appropriation -----	2
University Support -----	2
Institute Location -----	2
State Advisory Committee -----	3
RESEARCH PROGRAM -----	3
ANNUAL BASE GRANT (104 – B) -----	3
ND WRRI GRADUATE RESEARCH FELLOWSHIPS -----	6
PROJECTS (104-B: Fellowship)	
Uptake and effects of environmental estrogens on growth of fish----- <i>Fellow: Andrea Hanson</i>	8
Studies of Seasonal Succession of Cyanobacteria and Green algae at Heinrich-Martin Impoundment, North Dakota ----- <i>Fellow: Anusha Balangoda</i>	11
Source tracking of Cryptosporidium in rural watersheds ----- <i>Fellow: Brianna Schneck-Stenger</i>	14
Arsenic Contaminated Groundwater Remediation by Entrapped Nanoscale Zero-Valent Iron <i>Fellow: Christopher Capecchi</i>	16
Plant-based Biopolymers for Entrapping Metal Nanoparticles for Arsenic Removal: Biodegradation and Treatability Studies ----- <i>Fellow: Dhritikshama Roy</i>	18
Flood Risk Assessments of Various Scenarios for Devils Lake under GCM Downscaling Simulations Using a Coupled Hydro-Climatic Model Incorporating Recent Advances in Lake Evaporation Estimations. ----- <i>Fellow: Hasin Shahad Munna</i>	20
Integrating life stage habitat into landscape genetics model for the conservation of a declining amphibian species ----- <i>Fellow: Justin Fisher</i>	22
Assessment of Water Quality in Devils Lake using Satellite Imagery ----- <i>Fellow: Kate Overmoe-Kenninger</i>	24
Bioavailability of Dissolved and Colloidal Organic Carbon Bound Estrogen ----- <i>Fellow: Katrin Chambers</i>	25

Techniques of Assessing Changes in River Flooding Patterns -----	27
<i>Fellow: Kyle Haflinger</i>	
Ecosystem Services and Wetland Condition Assessment in the Prairie Pothole Region -----	28
<i>Fellow: Lindsey Malum</i>	
Hydrologic adaptation of SWAT model for snow dominated and high groundwater table conditioned watersheds and scenario analysis of impacts of tile drainage on stream flow -----	29
<i>Fellow: Mohammed Mizanur Rahman</i>	
Co-entrapment of iron nanoparticles and trichloroethylene degrading bacteria in alginate biopolymer for groundwater remediation -----	31
<i>Fellow: Sharnya Shanbhogue</i>	
Role of agricultural drainage on transport of Cryptosporidium oocysts in North Dakota -----	33
<i>Fellow: Tanush Wadhawan</i>	
Temporal-spatial distribution (dynamics) of phytoplankton and diversity in relation to lake physical and chemical condition -----	35
<i>Fellow: Veselina Valkov</i>	
NATIONAL COMPETITIVE PROGRAM (104 – G) -----	41
INFORMATION DISSEMINATION -----	38
PUBLICATIONS AND PRESENTATIONS -----	41
THESES AND DISSERTATIONS -----	43

## INTRODUCTION

This report describes the activities of the North Dakota Water Resources Research Institute (NDWRRI) during the period of March 1, 2011 to February 29, 2012.

The ND WRRI is one of the 54 institutes known collectively as the National Institutes for Water Resources (NIWR). The NDWRRI was founded in 1965, by authority of Congress (Water Resources Research Acts of 1964, 1972, 1984, and 1990), and is administrated through the United States Geological Survey. Section 104 of the Water Resources Research Act requires the NDWRRI to apply its Federal allotment funds to:

1. Plan, conduct or otherwise arrange for competent research that fosters: (A) the entry of new research scientists into the water resources field, (B) training and education of future water resources scientists, engineers, and technicians; (C) the preliminary exploration of new ideas that address water problems or expand understanding of water and water-related phenomena; and (D) the dissemination of research results to water managers and public.
2. Cooperate closely with other college and universities in the state that have demonstrated the capability for research, information dissemination and graduate training, in order to develop a statewide program designed to resolve State and regional water and related land problems.
3. Cooperate closely with other institutes and other organizations in the region to increase the effectiveness of the Institute and for the purpose of promoting regional cooperation.

This year, NDWRRI once again allocated its 104(B) resources to fund Graduate Fellowship research projects. The institute also continued its efforts to enhance communications between the State and Federal agency personnel and university faculty and students. NDWRRI also worked closely with the Environmental and Conservation Sciences program of North Dakota State University (NDSU), Natural Resources Management program of NDSU, and the International Water Institute, Fargo, ND on water related research issues and collaboration.

## **Program Management**

The Institute continued the same administrative mechanism with a director managing the institute program with the help of a State Advisory Committee. Dr. G. Padmanabhan, Professor of Civil Engineering, is the director. Linda Charlton, a NDSU employee, has been working part-time for the Institute to assist the director with Institute finances, communications and information transfer. The State Advisory Committee consists of three members representing the three principal water agencies in North Dakota: State Water Commission, State Department of Health, and the USGS North Dakota District. In addition, the Institute also seeks advice from the faculty of the two research universities of the State: North Dakota State University and University of North Dakota.

## **State Appropriation**

The State Water Commission continued its support of 15% match to the 2011 – 2012 the Graduate Research Fellowship program of NDWRRI under federal 104 (B) funding. This is eighth year the SWC provided support to the Fellowship program.

## **University Support**

North Dakota State University and the University of North Dakota administrations consider the NDWRRI activities important and are supportive of its efforts.

## **Institute Location**

The Institute continues to operate from the Administrative Building of the College of Engineering and Architecture of North Dakota State University in Fargo, North Dakota, The director may be reached at:

ND Water Resources Research Institute  
North Dakota State University  
Civil Engineering, Dept. # 2470  
Fargo, ND 58108-6050  
Phone: (701) 231-7043  
Fax: (701) 231-6185  
E-mail: G.Padmanabhan@ndsu.edu

## **State Advisory Committee**

The State Advisory Committee provided guidance on water resources research priorities in the State and region, and participated in the review and evaluation of research proposals and projects. The current committee members are:

Gregg Wiche, District Chief, U.S. Geological Survey, Water Resources Division,  
Bismarck, North Dakota

William Schuh, Water Appropriation Division, North Dakota State Water Commission,  
Bismarck North Dakota

Mike T. Sauer, Environmental Health Section, North Dakota Department of Health,  
Bismarck, North Dakota

The committee members are senior officials in the three major agencies in North Dakota responsible for much of the water resources research done outside of NDSU and UND in North Dakota.

## **RESEARCH PROGRAM**

### ***ANNUAL BASE GRANT (104-B)***

In the past several years NDWRRI has offered competitive fellowships to NDSU and UND graduate students for research on water resources topics under a Graduate Research Fellowship (GRF) program effectively using the modest amount of the 104(B) annual base grant. The program meets the requirements of Section 104 of the Water Resources Research Act of 1984. The fellowship program encourages entry of young university faculty and new research scientists into the water resources field; provides training and education to future water resource scientists and engineers; promotes exploration of new ideas that address water problems or expand understanding of water quantity, quality and related phenomena; and engages university faculty in collaborative research programs seeking supports from entities concerned with water problems.

This year, the NDWRRI continued the GRF program and applied bulk of the federal allotment to it. The GRF program is administrated and monitored by the director. Applications are invited from the graduate students and their advisors of the two research universities of the State, NDSU and UND. A rigorous review by the State Advisory Committee and other water professionals in the state determines the awards. Active participation of the academic advisors of the students in meeting matching requirement and seeking co-funding from local, state and other sources is another positive aspect of the program. Periodical review of the progress of the students in meeting the fellowship expectations is ensured by seeking reports from the students and by encouraging them to make presentations in local, regional, and national technical seminars and conferences.

Guidelines for the 2011-2012 Graduate Research Fellowship were posted on the Institute website in September 2010, and the request for applications was announced in the faculty news publications of the two university campuses in last week of October. The following is the request for application that was published on the UND and NDSU campus newsletters, and distributed by e-mail lists:

## **September 25, 2010 Issue of It's Happening at State (NDSU Publication)**

### **2011 ND WRRI graduate research fellowship applications invited**

The North Dakota Water Resources Research Institute (ND WRRI) has announced its 2011 Graduate Research Fellowship program.

North Dakota State University and University of North Dakota graduate students who are conducting or planning research in water resources may apply for fellowships varying from three summer months to a full year in duration. Typically in the past fellowship awards for master's degree students have been in the range \$800-\$1,000 and for doctoral students it has been \$1,000-\$1,400 per month. The fellowship funds must be applied between March 1, 2011, and Feb. 29, 2012.

Projects proposed for fellowship support should relate to water resources research issues in the state or region. Regional, state or local collaborations or co-funding will strengthen an application. Fellowships have a matching requirement of two non-federal dollars to one federal dollar. At the time of applying, applicants should have a plan of study filed and/or should have a thesis research topic selected. Applications need to be prepared in consultation with advisers. Advisers of the applicant should co-sign the applications. Applications from students and advisers who have not met the reporting requirements of their previous fellowship projects will not be considered for funding.

The general criteria used for proposal evaluation include scientific merit, originality of research, research related to state or region, and extent of regional, state or local collaboration and/or co-funding.

Applications are due in the office of the ND WRRI director by 5 p.m., Nov. 12. The proposals will be reviewed by a panel of faculty members and state water resources research professionals. Announcement of awards will be made by early January. Consult the ND WRRI Web site, [www.ndsu.edu/wrri](http://www.ndsu.edu/wrri), for background information on the program, and guidelines for preparation of applications. Applicants and advisers who are new to the program are urged to contact ND WRRI director, G. Padmanabhan, at 1-7043, or [G.Padmanabhan@ndsu.edu](mailto:G.Padmanabhan@ndsu.edu).

Send applications to Dr. G. Padmanabhan, Director of North Dakota Water Resources Research Institute, Civil Engineering, CIE 201E, NDSU Department 2470, P.O. Box 6050, Fargo, ND 58108-6050.



## **NDWRRRI GRADUATE RESEARCH FELLOWSHIPS**

Fellowships ranging from \$2000 to \$12,000 were awarded to nine graduate students from NDSU conducting research in water resources areas from 104(B) funding and three students from UND were awarded from ND State Water Commission support of the Fellowship program. NDSWC funding was also used for partially enhancing the Fellowship amounts of NDSU students. Selection of student Fellows and the award amounts were based on competitive proposals prepared by the students with the guidance of their advisers. Projects proposed for fellowship support should relate to water resources research issues in the state or region. Regional, state, or local collaboration or co-funding is encouraged. Fellowships have a matching requirement of two non-federal dollars to one federal dollar. A panel of state water resource professionals reviews the proposals and selects the Fellows and award amounts based on the quality of proposals and the priority of the proposed projects for the state and region. The general criteria used for proposal evaluation include: scientific merit, originality of research, research related to state and/or region, and extent of regional, state or local collaboration and/or co-funding.

This year, fifteen applications were received: eight Ph.D. and seven MS students. Highly competitive proposals and limited availability of funds restricted the amount of awards. Four of the Fellowships were renewals, three Ph.D and one MS. The renewals are Brianna Schneck, Anusha Balangoda, Dhritikshama Roy, and Andrea Hanson.

The titles of the fellowship projects awarded are given below and details are provided for each project under separate project sections.

## 2011-12 Fellows and their projects

2011-12 ND WRI Fellows, their advisers, and Fellowship research projects:

Andrea Hanson (Fellow), Biological Sciences; Mark Sheridan (adviser)  
Uptake and effects of environmental estrogens on growth of fish

Anusha Balangoda, Environmental and Conservation Sciences; Wei Lin  
Studies of Seasonal Succession of Cyanobacteria and Green algae at Heinrich-Martin  
Impoundment, North Dakota

Brianna Schneck, Biological Sciences; John McEvoy and Mark Clark  
Source tracking of *Cryptosporidium* in rural watersheds

Christopher Capecchi, Civil Engineering; Achintya Bezbaruah  
Arsenic Contaminated Groundwater Remediation by Entrapped Nanoscale Zero-Valent Iron

Dhritikshama Roy, Civil Engineering; Achintya Bezbaruah and Eakalak Khan  
Plant-based Biopolymers for Entrapping Metal Nanoparticles for Arsenic Removal:  
Biodegradation and Treatability Studies

Hasin Shahad Munna, Civil Engineering (UND); Howe Lim  
Flood Risk Assessments of Various Scenarios for Devils Lake under GCM Downscaling  
Simulations Using a Coupled Hydro-Climatic Model Incorporating Recent Advances in Lake  
Evaporation Estimations.

Justin Fisher, Biological Sciences; Craig Stockwell  
Integrating life stage habitat into landscape genetics model for the conservation of a declining  
amphibian species

Kate Overmoe-Kenninger, Earth Science System and Policy (UND); Xiaodong Zhang  
Assessment of Water Quality in Devils Lake using Satellite Imagery

Katrin Chambers, Soil Science; Francis Casey  
Bioavailability of Dissolved and Colloidal Organic Carbon Bound Estrogen

Kyle Hafliger, Civil Engineering (UND); Howe Lim  
Techniques of Assessing Changes in River Flooding Patterns

Lindsey Malum, Natural Resources Management; Edward Dekeyser and Jack Norland  
Ecosystem Services and Wetland Condition Assessment in the Prairie Pothole Region

Mohammed Mizanur Rahman, Ag. and Biosystems Engineering; Zhulu Lin  
Hydrologic adaptation of SWAT model for snow dominated and high groundwater table  
conditioned watersheds and scenario analysis of impacts of tile drainage on stream flow

Sharanya Shanbhogue, Civil Engineering; Achintya Bezbaruah and Eakalak Khan  
Co-entrapment of iron nanoparticles and trichloroethylene degrading bacteria in alginate  
biopolymer for groundwater remediation

Tanush Wadhawan, Civil Engineering; Eakalak Khan and John McEvoy  
Role of agricultural drainage on transport of Cryptosporidium oocysts in North Dakota

Veselina Valkov, Civil Engineering; Wei Lin  
Temporal-spatial distribution (dynamics) of phytoplankton and diversity in relation to lake  
physical and chemical condition

# Uptake and effects of environmental estrogens on growth of fish

**Project Number:** 2011SWC

**Fellow:** Andrea Hanson

**Adviser:** Mark Sheridan

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## Publication:

Hanson, A., Kittilson, J.D., McCormick, S.D., Sheridan, M.A., 2011. Effects of 17 $\beta$ -estradiol, 4-nonylphenol, and  $\beta$ -sitosterol on the growth hormone-insulin-like growth factor system and seawater adaptation of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture, in press*.

Hanson, A., Kittilson, J.D., Sheridan, M.A., 2011. Environmental estrogens inhibit growth of rainbow trout (*Oncorhynchus mykiss*) by modulating the growth-hormone insulin-like growth factor system. *In preparation*.

Hanson, A. and Sheridan, M.A. Effects of Environmental Estrogens on Organismal Growth and the Growth Hormone-Insulin-Like-Growth Factor System of Rainbow Trout. Annual Meeting of the Society for Integrative and Comparative Biology, Salt Lake City, UT, January 4-7, 2011.

Hanson, A. and Sheridan M.A. Environmental Estrogens Inhibit the Expression of Insulin-like Growth Factors 1 and 2 in the Liver and Gill of Rainbow Trout in vitro. Presented at the Annual Meeting of the Society for Integrative and Comparative Biology, Charleston, SC, January 3-7, 2012.

## Progress:

Rapid technological development over the last 60+ years has been accompanied by increased production, use, and disposal of an expanding array of chemicals. As a result, chemicals are introduced into the environment through the air, soil, and water. Increasing research shows the deleterious effects of environmental contamination on microbes, aquatic and terrestrial plants, invertebrates, fish, amphibians, birds, and mammals, including humans. Chemicals in the environment are known to have numerous toxicological effects, including disruption of the endocrine system of animals.

Of particular concern is a broad spectrum of natural and synthetic compounds that mimic estrogen. Environmental estrogens (EE) include endogenous and synthetic animal estrogens (e.g., 17 $\beta$ -estradiol), phytoestrogens ( $\beta$ -sitosterol), mycotoxins (e.g., zearalenone), organochlorine pesticides (e.g., DDT), polychlorinated biphenyls (PCBs), and alkylphenol polyethoxylates (APEs; e.g., 4-nonylphenol). EE enter soil and water through agricultural (including farm animal production) and veterinary (e.g., diethylstilbesterol, estradiol cypionate; widely used in dogs, cats, and horses) application, manufacturing, natural hydrological processes, and watershed/waste water management practices. Despite the fact that many EE

are found at levels below the legal limit, the potential long-term impact of chronic exposure to low levels is not clear, especially given that EE are rapidly transferred through the food web, accumulate in tissues, and can have epigenetic effects to affect progeny.

The goal of this project is to use rainbow trout as a model species to assess the biotic contribution to the fate of environmental estrogens in aquatic ecosystems by measuring their uptake and metabolism and by evaluating their impact on animal health as assessed by organismal growth. The hypothesis of this project is that environmental estrogens are taken up by rainbow trout from water and alter postembryonic growth by modulating the growth hormone (GH)-insulin-like growth factor-1 (IGF-1) system.

First, we have evaluated the effects of environmental estrogens on feeding and food conversion, and on postembryonic growth rate. In terms of overall organismal growth, the environmental estrogens, estradiol (E2),  $\beta$ -sitosterol ( $\beta$ S), and nonylphenol (NP) were to shown to lead to decreased organismal growth in terms of relative growth (both weight and length) compared to controls.

Next, we have completed work examining the effects of environmental estrogens on the GH-IGF system *in vivo*. All environmental estrogens were shown to decrease hepatic sensitivity to GH, as assessed by GH receptor 1 (GHR 1) mRNA expression. In addition to decreasing hepatic sensitivity, it was also determined that hepatic production of IGF-1 and IGF-2 decreased in environmental estrogen-treated fish. In gill, E2 reduced expression of mRNAs encoding GHR2, IGF-1, IGF-2, and IGFR1A, whereas NP reduced the expression of GHR 1, GHR 2, IGF1, IGF-2, and IGFR1A mRNAs.  $\beta$ S had no significant effect on the expression of any GH-IGF system component in gill filaments. The effects of EE on levels of GHR 1, GHR 2, IGFR1A and IGFR1B mRNA in red muscle and white muscle were somewhat variable.  $\beta$ S and NP decreased GHR 1 in red muscle, while GHR 2 mRNA expression decreased in white muscle. High concentrations of E2 reduced levels of IGFR1A in white muscle and of IGFR1B in red muscle, whereas NP reduced the expression of IGFR1A and IGFR1B mRNAs in red muscle.

These data support the hypothesis that EE reduce growth in fish by modulating the GH-IGF system in terms of GH sensitivity, IGF production, and IGF sensitivity in a tissue-specific manner at the molecular level and in terms of overall organismal growth.

Experiments were initiated to examine the effects of environmental estrogens on the GH-IGF system *in vitro* as assessed by mRNA expression of GHR1/2, IGF-1/2, and IGFR1A/B in liver or gill. E2,  $\beta$ S, and NP were not observed to affect the expression of GHRs in any tissues examined. However, decreased expression of both IGF-1 and IGF-2 in a time- and concentration-dependent manner was observed in the liver. Potency and efficacy of each of the environmental estrogens varied with E2 and NP eliciting the greatest response. Similar results were observed in gill. Experiments were conducted examining the *in vitro* response in muscle tissue of rainbow trout. These current data, in addition to the muscle data, will allow us to fully characterize the time course and efficacy of EE on IGF-1 binding characteristics and on the growth-promoting actions of IGF-1 in target tissues.

The findings of this project will advance our understanding of EE uptake and metabolism, and for the first time establish whether or not these agents act as endocrine

disruptors of growth in a wide-ranging aquatic species. While this work pertains specifically to rainbow trout, because the elements of the GH-IGF-1 system are highly conserved among vertebrates, the findings will have broad relevance to understanding the impact of EE in other aquatic and terrestrial species.

# **Studies of Seasonal Succession of Cyanobacteria and Green Algae at Heinrich-Martin Impoundment, ND**

**Project Number:** 2011ND237B

**Fellow:** Anusha Balangoda

**Adviser:** Wei Lin

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## **Publication:**

Anusha Balangoda (2011) Impact of Eutrophication on Water Quality at Heinrich-Martin Dam Impoundment, North Dakota”, Young Professionals **poster** competition at the North Dakota Water & Pollution Control Conference (NDWPCC), 11-13 October 2011, Bismarck

Anusha Balangoda (2011) “Population Variation of Cyanobacteria and Green Algae at Heinrich-Martin Dam Impoundment, ND”, Presented at the 5th International Student Prairie Conference, June 2-3, 2011. North Dakota State University, Fargo, ND.

Anusha Balangoda, Veselina Valkov, Jinhai Zhao, and Wei Lin (2012) “Seasonal Variations of Water Quality and Algal Growth in the Heinrich-Martin Dam Impoundment” Presented at the North Dakota Water Quality Monitoring Conference, February 27-29, Bismarck, North Dakota.

## **Progress:**

In North Dakota, major industry is agriculture including production of many crops, farms, and ranches. So the major sources of nutrients (Nitrogen and Phosphorous) loading into the ND's lake and reservoirs are erosion and runoff from animal feeding operations, and hydrologic modifications. So this excessive level of phosphorous and nitrogen may result in an overstimulation of nuisance algal blooms. According to the seasonal variation when lakes become more eutrophic, with the increase of phytoplankton biomass the diversity of the assemblage decreases ultimately leading to the dominance of cyanobacteria (Blue green algae). Blue-green algal blooms (Harmful Algal Blooms-HABs/Cyanobacteria blooms) produce toxins are poisonous to animal and human health in waters used for recreational and drinking purposes. So in contrast to other phytoplankton, cyanobacteria can withstand under wide range of environmental conditions which is related to their specific growth characteristic features. So eutrophication is not only being a state wise problem it is one of the most widespread environmental problems of inland waters in the world.

The main objective of this study is to identify population variation of cyanobacteria and green algae in relation to their growth requirements.

## Impacts of Aeration on the Water Temperature & the Dissolved Oxygen (DO)

Seasonal variations of water temperature and DO were observed in 2010 and 2011. In addition to seasonal variation, thermal stratification was observed after aeration was turned off for a month in 2011. Thermal stratification was established only in the deeper part of the impoundment after aeration was turned off. When the aeration was in operation the lake showed a mixed condition. The water temperature difference from surface to bottom of the water column was less than 0.5°C (Fig. 1a). DO also showed pronounced vertical variation along the depth. A rapid DO drop from surface (7 mg/L) to 2 m (1 mg/L) was observed after aeration was turned off. The DO concentration less than 1 mg/L was observed near the bottom in the deeper part of the impoundment over the stratified period.

## Impacts of Aeration on Nutrient Concentrations

Nitrogen (N) and phosphorous (P) are the nutrients that limit plant and algae growth. TDIN ( $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and  $\text{NO}_2^-$ ) and SRP (soluble reactive phosphorous) are the available forms for algal growth. Both TDIN and SRP showed seasonal variations during sampling in 2010 and 2011. In addition to seasonal variation, variations of nutrients concentration with the condition of aeration and after the aeration was turned off were also observed. Nutrients were mostly distributed from surface to bottom in the lake when the aeration was in operation. However, when the aeration was turned off, nutrients concentration varied along the depths in deeper part of the impoundment. The SRP concentration showed concentration gradient from surface (0.06 mg/L) to bottom (0.45 mg/L) may indicate release of phosphorous from sediment to overlying water column (Fig. 2a). Similar to SRP, the TDIN concentration also showed concentration gradient from surface (0.08 mg/L) to bottom (1.1 mg/L). The TDIN concentration was dominated by  $\text{NH}_3\text{-N}$  near the bottom (1.08 mg/L) shows that possible release of  $\text{NH}_3\text{-N}$  from sediment to water column. Further, results showed that N and P concentration gradient from surface to bottom was significant in HMD over the stratified period. Therefore, the mechanisms and factors that affecting release of nitrogen and phosphorus from sediments to water column is needed to be studied in HMD.

The TDIN: SRP ratio varied seasonally as well as under the condition of aeration and after the aeration was turned off during sampling in 2010 and 2011. The TDIN: SRP ratios changed from 6.44:1 (with aeration) to below 2:1 (without aeration), which was less than the cellular element mass ratio (7.2:1) of algae, indicates N limiting condition in HMD (Fig. 2c). Therefore, the factors that affecting for N limitation is needed to be determined in HMD. Further, results showed that total phosphorous (TP) concentration was  $\geq 100 \mu\text{g/L}$  in HMD. Limnologist and lake managers predict nuisance algal blooms with TP concentration greater than about 0.01 mg/L (Gibson et al., 2000). Therefore, determination of what causes P rich condition in HMD is also needed.



### Impacts of Aeration on the Chlorophyll a (Chl a) Concentration

Chl a is the primary photosynthetic pigment, so a measure of its concentration in a water sample is represent total algal biomass. Chl a showed a seasonal variation over the study period in 2010 and 2011. Under the condition of aeration, Chl a concentration was mostly homogenized throughout the water column. However, there was a significant vertical variation of Chl a was observed in deeper part of the impoundment after the aeration was turned off about a month in 2011. Chl a showed a high concentration ( $>80 \mu\text{g/L}$ ) from surface/Secchi depth level to a low concentration ( $<10 \mu\text{g/L}$ ) near the bottom over stratified period. The seasonal stratification of water columns determines the general availability of the resources light and nutrients for phytoplankton growth. Results showed that N and P were accumulated near bottom. Therefore, nutrients may not have adequate supply to a well-mixed top layer after the aeration was turned off. So, what determines their vertical distribution in stratified water columns in HMD are needed to be studied. In addition to Chl a, Secchi depth also measures trophic status in lakes. The Secchi depth showed a seasonal variation over the study period and Secchi depth was  $\geq 1$  m over most of the sampling dates. The lowest Secchi depth (0.7 m) was observed after the aeration was turned off.

### Impacts of Aeration on the Phytoplankton Population

Seasonal variation of phytoplankton population shift was observed during sampling in 2010 and 2011. In addition to seasonal variation, phytoplankton population shift under the condition of aeration and after the aeration was turned off also observed. Algal bloom condition ( $2.88 \times 10^6$  Cells L<sup>-1</sup>) was observed in HMD when aeration was in operation. Surface water showed green color balls like aggregates and those were identified as *Aphanizomenon* sp. clusters under inverted microscope analysis. After a week when aeration was turned off, large balls like aggregates showed more dispersed condition and concentrated into the surface layer. A significant decrease of cyanobacteria cell densities (from  $4.58 \times 10^6$  cells L<sup>-1</sup> to  $1.14 \times 10^5$  Cells L<sup>-1</sup>) were observed at the surface layer over the stratified period. A Cyanobacteria population, which represented much of the total phytoplankton population, was dropped significantly after the aeration was turned off.

# Source Tracking of *Cryptosporidium* in Rural Watersheds

**Project Number:** 2009ND183B

**Fellow:** Brianna Schneck-Stenger

**Adviser:** John McEvoy and Mark Clark

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## Progress:

The Upper Midwest has among the highest incidences of human cryptosporidiosis in the U.S. There is therefore a critical need to determine *Cryptosporidium* sources and transmission dynamics in this region. A study, which I co-authored, has shown that most human cryptosporidiosis in the region is caused by *C. parvum*, a species associated with humans and cattle. However, the traditional model, describing *Cryptosporidium* movement from livestock to humans via water, appears over simplistic in light of recent evidence of wildlife sources of contamination. To better understand the human health significance of *Cryptosporidium* in rural watersheds, we need to determine the source of the contamination.

The objective is to determine the factors influencing the contributions of cattle and wildlife to *Cryptosporidium* in rivers. This is critical to understand the public health significance of *Cryptosporidium* in rural watersheds.

Empirical evidence supports host adaptation and a limited host range for most *Cryptosporidium* species and genotypes. This knowledge has been used to track sources of water contamination and characterize human health risk; however, the factors affecting host range remain unclear. We are using small mammals (mainly rodent species) as a model to study the effect of factors such as host population density, diversity and behavior on the host range of *Cryptosporidium* genotypes.

Spring flooding results in the movement of *Cryptosporidium* from the fields of North Dakota, South Dakota and Minnesota to the Red River. Livestock contributed significantly to *Cryptosporidium* contamination in the Red River during major spring floods. Cattle were the primary source of surface water contamination. We can estimate the flow of oocysts during peak flooding in 2010 at 728,000 per second, based on our oocyst counts and a river flow of 560,000L per second. That is significant when it can take less than 100 oocysts to infect a host.

Small mammals are hosts to various genotypes of *Cryptosporidium*. We found *Cryptosporidium* species or genotypes in over 42% of our samples. We now have more

than 600 samples and the prevalence is approximately 40%. We've identified at least 14 *Cryptosporidium* genotypes/species in 10 wildlife host species. We have found a novel genotype of *Cryptosporidium* in Eastern chipmunks and are planning to publish the findings once further molecular characterization and microscopic analysis is complete. Two species of *Cryptosporidium*, *C. parvum* and *C. ubiquitum*, are considered human pathogens, and were found in a number of wildlife samples.

# **Arsenic Contaminated Groundwater Remediation by Entrapped Nanoscale Zero-Valent Iron**

**Project Number:** 2011ND238B

**Fellow:** Christopher Capecchi (work completed by a few other graduate and undergraduate students since Mr. Capecchi quit the Fellowship half way through)

**Adviser:** Achintya Bezbaruah

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## **Publication:**

Capecchi, C.; Bezbaruah, A. Novel Alginate Entrapped Nanoparticles for Groundwater Arsenic Remediation, Proc. World Environmental and Water Resources Congress 2011, pp. 3389-3395, 2011 (also oral presentation).

## **Progress:**

In response to the significant threat arsenic presents, the United States Environmental Protection Agency (USEPA) drastically modified the maximum contaminant level (MCL) for arsenic in drinking water from 50  $\mu\text{g/L}$  to 10  $\mu\text{g/L}$  in 2006. Even at 10  $\mu\text{g/L}$  the world health organization (WHO) estimates a 0.2% chance of developing cancer in humans. Millions are presently at risk due to high arsenic levels in drinking water. Acute and chronic arsenic exposure from drinking water has been reported in many countries, most of which have large proportions of drinking water contaminated with high concentrations of arsenic (total As 50  $\mu\text{g/L}$ ). The United States Geological Survey (USGS) analysis of 30,000 random groundwater samples across the United States found that approximately 10% of sites had arsenic concentrations in excess of 10  $\mu\text{g/L}$ . Moreover, arsenic is second only to lead (Pb) as the most commonly found hazardous contaminant at Superfund Sites. In Southeastern North Dakota more than 25% of groundwater samples contain arsenic at levels in excess of 10  $\mu\text{g/L}$ . The U.S. EPA conducted a five year review (2004-2008) to analyze the remedial action implemented in Southeastern North Dakota. Approximately 375 wells in the 26 townships (about 568 square miles) were sampled and it was found that more than 84% samples have the arsenic concentration above the MCL. This presents an unacceptable risk of cancer and adverse health effects to the residents who depend upon water from the aquifer. By entrapping NZVI within a biopolymer (Ca-alginate) the overall contact time with contaminants will be prolonged, allowing individual particles to react more efficiently. By optimizing critical parameters such as NZVI dosage and pH, the waters can be effectively remediated below the USEPA arsenic MCL (10  $\mu\text{g/L}$ ). The objective of this research is to determine the treatability of arsenic by entrapped NZVI. The specific objectives of the study are: 1) Conduct entrapped NZVI treatability batch studies with various  $\text{As}^{5+}$  and  $\text{As}^{3+}$  concentrations. 2) Examine the effects of individual ions (which are present in groundwater) on the arsenic removal (by entrapped NZVI) reaction kinetics. 3) Characterize entrapped NZVI within Ca-alginate beads using X-ray

diffraction (XRD) and scanning electron microscopy (SEM) to understand the arsenic treatment mechanisms. 4) Perform entrapped NZVI treatability batch studies with actual arsenic contaminated groundwater.

This project explored aqueous arsenic removal using nanoscale zero-valent iron (NZVI) entrapped in calcium (Ca) alginate beads. The results from this study show great promise for entrapment technique as an advanced treatment technique for aqueous arsenic. Arsenic is a serious threat to human health and millions of people are affected by arsenic contamination in various parts of the world including the United States. The entrapment process reduces mobility of the nanoparticles by confining them within the polymer matrix and, thus, reducing the risk of post-treatment hazard by arsenic sorbed onto NZVI. Ca-alginate polymer is an excellent choice as an entrapment medium as it is non-toxic and has little solubility in water. In bench scale batch studies with initial As(V) concentrations of 1-10 mgL<sup>-1</sup>, ~ 80-100% arsenic removal was achieved within 2 hours. While the reaction kinetics differ between bare and entrapped NZVI, the overall reductions of arsenic are comparable. Surface normalized arsenic reduction reaction rate constants ( $k_{sa}$ ) for bare and entrapped NZVI were 3.40-5.96x10<sup>-3</sup> and 1.90-4.43 x10<sup>-3</sup> L m<sup>-2</sup> min<sup>-1</sup>, respectively.

Graduate student who originally worked on this project was Christopher Capecchi, and later two other graduate students and a few undergraduate students participated in this project. One peer reviewed paper (2011) has been published in the Proceedings of ASCE-EWRI World Environmental and Water Resources Congress 2011, and another paper has been submitted (2012) to a referred journal for possible publication. In addition two posters were presented on the research (both in 2010).

# **Plant-based Biopolymers for Entrapping Metal Nanoparticles for Arsenic Removal: Biodegradation and Treatability Studies**

**Project Number:** 2011ND239B

**Fellow:** Dhritikshama Roy (Work completed by a former NDWRRI Fellow, Rabiya Shabnam, and two undergraduate students from Civil Engineering since Ms. Roy quit the Fellowship early on practically with no contribution to the project)

**Adviser:** Achintya Bezbaruah

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## **Publication:**

Roy, D.; Kalita, H.; Khan, E.; Bezbaruah, A. Fate of Polymeric Nanoparticle Delivery Vehicles in the Aquifer: Biodegradation of Polydimethylsiloxane, 2011 World Environmental & Water Resources Congress, Environmental & Water Resources Institute, American Society of Civil Engineers, Palm Springs, CA, May 2011.

## **Progress:**

Nanoscale zero-valent iron (nZVI) particles have been attractive for remediation of various contaminants including arsenic. However, because of the smaller particle size and relatively higher dispersibility, nZVI becomes mobile in aquifers. Moreover, in higher concentration, nZVI tend to agglomerate due to magnetic and Van der Waals forces. This allows them to form large particle and settle into aquifer media pores. Agglomerated particle has decreased specific surface and hence lose the reactivity that individual nZVI has. Polymers are used to coat or entrap nZVI to overcome the problem of agglomeration or uncontrolled dispersion, Such entrapped nZVI have shown higher reactivity towards contaminants. However, polymers synthesized in previous research are not biodegradable or shows limited biodegradation. Lack of biodegradation may limit the use of these polymers in groundwater where polymers may themselves become pollutants. It is imperative to development and use polymers which are easily biodegradable to benign end product in the aquifer. This project proposes to use plant-based (bio) polymers to entrap nZVI and study microbial biodegradation of the polymers used. A soybean-based biopolymer will be the primary focus of this study for removal of Arsenic.

Nanoparticles (mostly iron-based) have been successfully used for groundwater contaminant removal. However, they are agglomerated in aqueous media due to magnetic and Van der Waals forces and settle down in the aquifer pores. Surface modification of nanoparticles have been tried using synthetic and plant-based biopolymer within this NDWRRI supported project on “Plant-based Biopolymers for Entrapping Metal Nanoparticles for Arsenic removal: Biodegradation and Treatability Studies”. The project was originally started by a graduate student from Environmental Conservation Sciences and later continued by a former NDWRRI Fellow (Rabiya Shabnam) and two undergraduate students from Civil Engineering under the title “Testing Biodegradation of

Polymer Coated Nanoparticles in Aqueous Media”. Biodegradation of synthetic and plant-based polymers in the presence of nanoscale zero-valent iron (NZVI) has been the emphasis of this project. The copolymers developed for NZVI coating by two former NDWRRI fellows (Sita Krajanpan and Harjyoti Kalita) were used in the degradation studies. The results indicate that polysiloxane-polyethylene glycol-carboxylic acid based synthetic copolymers do not degrade easily but degrade faster in the presence of NZVI. A soybean oil based copolymer used in this study showed complete and fast degradation with and without NZVI. This fellowship research has used a new robust method to evaluate biodegradability of polymers and polymer coated NZVI. The new method uses respirometric techniques and acquires biochemical oxygen demand data continuously from the aqueous samples containing polymer or polymer coated NZVI.

# **Flood Risk Assessments of Various Scenarios for Devils Lake under GCM Downscaling Simulations Using a Coupled Hydro-Climatic Model Incorporating Recent Advances in Lake Evaporation Estimations.**

**Project Number:** 2011SWC

**Fellow:** Hasin Shahad Munna

**Adviser:** Howe Lim

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## **Progress:**

Located in the Ramsey and Benson County, Devils Lake is a closed basin that receives surface runoff from 3,810 square miles of drainage basin (USGS). The uncontrolled growth of the lake has been an alarming issue for North Dakota for the last few years. Over the past 15 years, the North Dakota State Water Commission (NDSWC), in collaboration with the Devils Lake Basin Joint Water Resource Board, has implemented numerous flood mitigation projects for Devils Lake, costing more than \$500 million, to raise levees, improve roads, and build an outlet. The rising water has destroyed hundreds of homes and businesses, inundated thousands of acres of productive farmland, and displaced nearly 1,000 people, about 15% of the population. The on-going lake flooding is both unique and complex which warrants much greater research efforts especially in the water resources engineering and management perspectives.

The proposed research consists of 3 basic parts.

Modification and recalibration of different models to estimate the evaporation time series: A total of eight radiation based evaporation models have been analyzed to verify the best model in Devils Lake region. Among these the Hargreaves method, Doorenbos and Pruitt method and Makkink method showed relatively close matches with the observed values from nearby gauging stations. A simplified Penman equation was also examined to check its efficiency in estimating the evaporation. The research will focus on recalibrating the proposed methods to obtain better estimations.

Applying the simulated GCM based temperature and precipitation time series to the calibrated and coupled hydro-climatic model to generate future lake levels: In this part the generated synthetic daily temperature and precipitation time series will be applied to the coupled model. Part of the task is to identify a common compatible format to import the synthetic datasets of numerous different scenarios into the coupled model. The HEC model uses a powerful database tool called HEC-DSSVue to import time series parameters. DSSVue can import data in ASCII format which can be converted into .dss format by using utility plug-ins. A properly oriented ASCII format will be designated for the interchange of data. After identifying the proper way for exchanging data, simulations



will be run which will give us future simulated water levels of Devils Lake at different selected time periods (2020, 2050) under varying climatic conditions.

Frequency analysis of the synthetic peak stages of Devils Lake:

There are different methods for fitting the probability distribution of peak values some of which are 2-parameter lognormal (LN2), 3-parameter lognormal (LN3), Gumbel distribution, general extreme value (GEV), Log-Pearson 3 (LP3) and Pareto distributions. For estimating PD for peak flood frequency analysis, the Log-Pearson Type III (LP3) distribution is recommended (Bulletin 17B) according to the U.S. Water Advisory Committee on Water Data (1982). Peak stage analysis will be done using the recommended method and also by other popular methods which will give a comparative measure of the goodness of fit. The distribution method with the best fit and also the recommended LP3 method will be applied in the frequency analysis of the simulated data to estimate peak stages for different return periods.

A hydro climatic model has been developed to estimate the lake behavior for a 20 to 50 year time period. Several sets of modeling software and heterogeneous data have been used in accomplishing our research goal. The watersheds that feed directly into Devils Lake were modeled using Arc-GIS and imported into HEC HMS model (watershed modeling software developed by the U.S. Army Corps of Engineers) to simulate the run-off to the Devils Lake which was calibrated by observed storm events. Then the run-off time series was imported into HEC-ResSim model to estimate the lake levels which was also calibrated with observed elevations. The calibration was done for the years of 2001 to 2004. The combined physical properties (Elevation-Storage-Area) were used to model the lake as during our study period the elevation of the lake was consistently over 1447 ft amsl. Now we are working on combining all the individual run-off sub-models into a single model which will then be coupled with the calibrated HEC-ResSim to permit simulations of the future time periods.

# **Integrating life stage habitat into landscape genetics model for the conservation of a declining amphibian species**

Project Number: 2011SWC  
Fellow: Justin Fisher  
Adviser: Craig Stockwell  
Start Date: 3/1/2011  
End Date: 2/29/2012

## **Publication:**

### **Presentations:**

Fisher, J., K. Purcell and C. Stockwell. Initial Survey of the Genetic Diversity of Northern Leopard Frog Populations in North Dakota. North Dakota Chapter of the Wildlife Society Annual Meeting, Mandan, ND, 02/2011.

Fisher, J., K. Purcell and C. Stockwell. Survey of the Genetic Diversity of Northern Leopard Frog Populations in North Dakota. Joint Meeting of Ichthyologists and Herpetologists, Minneapolis, MN, 07/2011

Fisher, J., K. Purcell and C. Stockwell. Updated Survey of the Genetic Diversity of Northern Leopard Frog Populations in North Dakota. NDSU ECS Graduate Student Poster Competition, Fargo, ND, 10/2011

## **Progress:**

What was once thought to be a highly abundant amphibian throughout much of North America, the northern leopard frog (*Rana pipiens*) has been recently petitioned for federal protection. Research and monitoring programs indicate widespread population declines and the U.S. Fish & Wildlife Service is currently seeking species information, particularly on genetic health. The northern leopard frog is unique compared to other amphibians in North Dakota in the fact that it behaviorally evades harsh winter weather by over-wintering in available wetlands at the water/mud interface. However, this over-wintering strategy can be deadly if freezing temperatures (i.e., ice) extend to the wetland bottom. Therefore, northern leopard frogs require wetlands with deep water, to survive winter. However, recently reported that during severe drought events in North Dakota, such drought refugia become uncommon. Thus, the spatial and temporal distribution of drought refugia can play an important role in the distribution of northern leopard frogs on the landscape. Accordingly, understanding where populations persist during past climate extremes can be used to provide insights into the role of drought refugia in northern leopard frog population dynamics, particularly in terms of genetic diversity and gene flow across the landscape.

Objectives are:

1. Examine population genetic structures at drought refugia sites to understand genetic sources for potential re-colonization of northern leopard frog populations in ND
2. Develop landscape genetic models to evaluate current influences of gene flow on northern leopard frog populations in ND

Field sampling and about 60% of the laboratory work have been completed. Baseline data analyses has been completed as well. My future work aims to increase the number of microsatellite markers to a total of 10 markers and examine DNA from another 20 populations. I anticipate an additional field sampling season (summer 2012) to gather additional populations to further evaluate my observed geographic distribution of genetic diversity and to also evaluate how genetic structure is correlated with various natural and anthropogenic landscape features.

# Assessment of Water Quality in Devils Lake using Satellite Imagery

Project Number: 2011SWC

Fellow: Kate Overmoe-Kenninger,

Adviser: Xiaodong Zhang

Start Date: 3/1/2011

End Date: 2/29/2012

## **Publication:**

Overmoe, K., X. Jia, W. Lin, & B. Kratz. Submitted October 2011. Evaluation of an Aeration System to Increase Dissolved Oxygen Level at an Impoundment in the Northern Great Plains. Water Research.

Gerla, P., M. Dinger, E. McCoy, K. Overmoe, & N. Roehrdanz. 2011. Monitoring and Modeling Long-Term Changes of Rainfall-Runoff Response and Water Quality in a Small Agricultural Watershed Restored to Prairie. The Geological Society of America.

Hill, M., S. Fore, K. Overmoe, Q. Zhou, R. Lemmons, & C. New. 2011. Dynamics, Metrics, Services and Scenarios for Ecosystems in North Dakota and Northern Minnesota, USA. NASA Carbon Cycle & Ecosystems Joint Science Workshop. (poster).

Overmoe, K. 2011. Comparison of Two Spatial Interpolation Methods for Secchi Disk Transparency in the Devils Lake Basin, ND. Advanced Geographic Information Systems Poster Session, UND. (poster).

Overmoe, K. 2011. Assessment of Water Quality in North Dakota Lakes and Reservoirs using Satellite Imagery. Earth System Science & Policy Seminar.

Overmoe, K. & A. Hewitt. 2011. Tree Extraction from LiDAR. Earth System Science & Policy

## **Progress:**

Historical water quality data (SDT, chl-a, total P, and turbidity) and data locations within all North Dakota lakes and reservoirs have been obtained from the EPA STORET, EPA Legacy, and USGS databases. Corresponding Landsat scenes and acquisition dates have been identified and image processing is currently in progress. In addition, I am currently working on the initial efforts of a manuscript analyzing long-term trends in water clarity in North Dakota lakes and reservoirs with special focus on lakes with historic (1970s) to current water quality data sets. These lakes include Ashtabula, Audobon, Darling, Devils, Sakakawea, and Tschida. The first field season in the Devils Lake basin was completed in 2011, with a total of six sampling dates corresponding with Landsat overpass from June to October. In addition, this was the first season deployment of a water quality monitoring buoy in Stump Lake. The buoy measured, recorded, and transmitted data on salinity, transparency, total dissolved solids, oxygen concentration and chl-a concentration every 10 minutes for 77 days. Collaboration efforts with the NDDOH and Prairie Waters Education and Research Center have begun to initiate a citizen volunteer effort for the summer of 2012. Additional contacts will be made with North Dakota high school teachers at the North Dakota Science Teachers Association conference held in February 2012 to increase participation in this study.

# Bioavailability of Dissolved and Colloidal Organic Carbon Bound Estrogen

Project Number: 2011ND242B  
Fellow: Katrina Chambers  
Adviser: Francis Xavier McKeon Casey  
Start Date: 3/1/2011  
End Date: 2/29/2012

## Progress:

The natural estrogen, 17 $\beta$ -Estradiol (E2), is the most potent endocrine disrupting compound. The issue of reproductive hormones in the environment is of particular relevance to animal agriculture because of the association of hormone detections with manure management practices. Natural hormone concentrations have been detected in runoff and receiving surface waters as a result of field manure application. Reproductive hormones have also been measured in subsurface waters in proximity to intensive livestock production. Widespread E2 detections throughout the soil profile and shallow groundwater in fields in and around a swine (*Sus scrofa domesticus*) farm has been found in western North Dakota.

The objectives of this proposed project are the following: (i) Evaluate the association of radiolabeled estradiol with both the Dissolved organic carbon (DOC) and colloidal organic carbon (COC) fraction derived from liquid swine manure, soil and liquid swine manure applied to soil (ii) Simulate rainfall events to investigate the dislodgment of radiolabeled estradiol from the COC fraction (iii) Evaluate the estrogen potency of estradiol bound to the DOC/COC, DOC and COC fraction using estrogen receptor competitive assays.

1. Preservatives: A preliminary survey of ELISA responses to various preservatives, E2 concentrations, and DOC/COC concentrations has been conducted. There is a potential that the preservative formaldehyde could interfere with the ELISA response. ELISA responses were measured for solutions of formaldehyde (2.5%) and another preservative, sodium azide (0.2%). Also, several concentrations of E2 and dilutions of DOC/COC solution were tested. Appropriate replications and blocks from this first ELISA analysis allowed the use of an ANOVA statistical comparison. The effects from this preliminary test indicated that COC/DOC (probability (p) = 0.0055), formaldehyde (p= 0.0420) and E2 concentration (p<0.0001) were significant in explaining the ELISA response. The conclusions that can be made from these initial results are: (i) Use sodium azide to preserve samples, (ii) The DOC/COC does have a significant effect on spectrometer response of the ELISA (i.e. the more DOC/COC, the greater the overestimation of concentration). However, the 5% and 0.5% COC/DOC concentrations were not significantly different from the solution with no DOC/COC. This may indicate that “yes” the E2 bound to DOC/COC can still interact with the hormone receptors and induce an estrogenic response.

2. E2 Association/Dissociation with COC: The results from the ultrafiltration of the swine lagoon manure spiked with radiolabeled E2 show that E2 is associated with COC particles and that E2 can be dislodged from the COC particles when rinsed with water. The environmental implication of these findings is that E2 can “hitch-hike” on COC particles to be distributed in the environment and be dislodged potentially as free estradiol.

3. E2 Bound to COC is Characteristic of a Particulate: Reverse and normal phase Thin Layer Chromatography (TLC) has been done on the COC fraction of liquid swine manure. The results show that the radiolabeled E2 bound associated with the COC fraction does not behave like a molecule but more as a particle. Specifically, the COC fraction did not respond to chromatographic separation because the radioactivity stayed at the origin of both the normal and reverse phase TLC. This indicates that the <sup>14</sup>C-E2 associated with the COC is acting like a particle and not like a molecule.

4. E2 Bound to COC is Bioavailable: The results from the Estrogen Receptor (ER) competitive-binding assays show that there is an estrogen response induced by the E2 associated with the COC fraction. The concentrations above the bars in Fig. 5 are the estradiol equivalence derived from the standard curve. These results indicated that estradiol bound to COC still has endocrine disrupting properties and can potentially cause adverse effects to aquatic organisms. The estrogen response from the COC rinses mimics the response of the radiolabeled estradiol experiments. This indicates that estradiol can be carried by COC particles but is loosely bound and can be released into the environment as free estradiol.

# Techniques of Assessing Changes in River Flooding Patterns

Project Number: 2011SWC

Fellow: Kyle Hafliger

Adviser: Howe Lim

Start Date: 3/1/2011

End Date: 2/29/2012

## Progress:

The purpose of this research is to study changes in the flooding patterns of rivers or streams in the Upper Midwest. Some rivers or streams have a significant increase or decrease in the mean annual peak flows in the past forty years than it did in the forty year or longer period prior to that. All of the river flow data used in this research came from the United States Geological Survey (USGS) website. This website has data on various statistics of river data, such as flow rates in this case. Annual and daily flow values were used in the research.

Eight different rivers were chosen to analyze the annual peak flows during the period of record. In the Midwest, the James River had a significant increase in the average annual peak flow since 1970. The time period from 1971 to present has over twice the value for the average annual peak flow compared to the time span of 1929 through 1970, a bigger change than any other river analyzed to find changes. Therefore, the James River Basin was chosen for study. The method used to find the magnitude of change in the mean annual peak flow during two specific time periods is the two tailed t-test. The two time periods used was from the period prior to 1970 and the period from 1971 to present. The rivers or streams that were chosen for analyzing were ones that had a P value of less than 0.05 from the two tailed t-test result. The lower the value of P, the bigger change that has occurred in the peak annual flows. Of the rivers researched in the Upper Midwest, only two had P values under 0.05: the Thief River at Thief River Falls, Minnesota, and the James River at Scotland, South Dakota. The James River had the lowest P value.

The daily flow data was analyzed for each of the rivers that had a significant change in the annual peak flows to look at the flood changes in more detail. A wavelet analysis using MATLAB of daily flows in the James River, at the Scotland, South Dakota station, showed an increase in flooding frequency starting around 1960. Most prominent frequency was found to be of one year.

The statistics of the temperature and precipitation haven not been analyzed yet. Other parameters will also be looked at to see what all factors contribute to the increase in the flooding. Several sites in the river basin will be analyzed for the causes of flooding changes.

# **Ecosystem Services and Wetland Condition Assessment in the Prairie Pothole Region**

Project Number: 2011ND241B

Fellow: Lindsey Meyers Malum

Adviser: Edward Dekeyser and Jack Norland

Start Date: 3/1/2011

End Date: 2/29/2012

## **Publication:**

Meyers, L.M., E.S. DeKeyser, J.E. Norland, C.L.M. Hargiss, and T. DeSutter. 2012. Wetland assessment and ecosystem services. North Dakota Water Quality Monitoring Conference, Bismarck, ND.

DeKeyser, E.S., C.L.M. Hargiss, L.M. Meyers, M.J. Ell. 2012. The development of a multiple collaborator venture connected to the National Wetland Condition Assessment in North Dakota. North Dakota Water Quality Monitoring Conference, Bismarck, ND.

## **Progress:**

This study seeks to assess and evaluate wetlands across the state of North Dakota with a special emphasis on wetlands in the Prairie Pothole Region (PPR). In the summer of 2011, four assessment and nutrient studies were completed on 55 wetlands. The National Wetland Condition Assessment (NWCA) evaluated vegetative, soil, water, algal, hydrological and buffer wetland characteristics. The NWCA not only included intensive sampling of these biological and physical criteria, but also included a rapid assessment of these criteria. Regional wetland assessments developed for North Dakota were also completed at each site. Each wetland was rapidly assessed using the North Dakota Rapid Assessment Method (NDRAM), plant community composition of each wetland was evaluated using the Index of Plant Community Integrity (IPCI), and functional characteristics of the wetlands were evaluated using the Hydrogeomorphic (HGM) model. At each wetland, live plant and soil samples were collected for nutrient analysis.

The final completion NDWRRI report has been submitted and is available on the NDWRRI website.



# **Hydrologic adaptation of SWAT model for snow dominated and high groundwater table conditioned watersheds and scenario analysis of impacts of tile drainage on stream flow**

Project Number: 2011SWC

Fellow: Mohammed Mizanur Rahman

Adviser: Zhulu Lin

Start Date: 3/1/2011

End Date: 2/29/2012

## **Publication:**

- Lin, Z., A.P. Kirilenko, and M.M. Rahman (2011). Coping with uncertainty in assessing climate change impacts on streamflows. Proceedings of the AWRA 2011 Spring Specialty Conference, Baltimore, MD.
- Rahman, M.M., and Z. Lin (2011). Hydrologic adaptation of SWAT model to analyze impacts of tile drainage on streamflows in snow dominated upper Red River of the North Basin. ASABE Annual International Meeting 2011, Louisville, KY.

## **Progress:**

USGS (U.S. Geological Survey) eight digit coded watersheds within the Red River basin of the North, USA are quite different from others for various reasons.

Firstly, this basin crosses three snow dominated states North Dakota, South Dakota and Minnesota where a greater part of annual precipitation comes from snowmelting in the late spring and early summer, on the other hand, throughout the summer and in early fall major contribution of runoff to streams is from rainfall. So, parameterization of any continuous model like SWAT (Soil and Water Assessment Tool) in such a mixed environment would be a difficult task. Secondly, intensive and diversified cropping practices in almost all watersheds not only make this area very uneven regarding water balance spatially and temporally but also contaminate soil and water (surface and subsurface) mass through non-point source of inorganic agricultural chemicals like fertilizers, insecticides, pesticides, herbicides etc. On average, the groundwater (GW) table fluctuates within 4 to 8 feet bgl (below ground level) and due to having comparatively high GW table, groundwater resource is subjected to be contaminated by various point & non-point sources, mainly from agricultural chemicals. Tile drainage has been becoming more popular in this region from few years back. This recent rapid trend of adopting tile drainage practices by farmers is making regional watershed's hydrological behaviors somewhat different than that of usual watersheds.

The aforementioned issues should be addressed during parameterization of the SWAT model developed by USDA-ARS (Arnold et al. 1993) that would be helpful for future

planning and managements. No specific research has been done in the past to parameterize the SWAT model for the impact analysis of tile drainage in the Red River basin of the North.

The specific objectives of this study are:

1. To optimize the hydrologic parameters of SWAT model for snow dominated watersheds of North Dakota
2. To identify the most sensitive parameters that must be paid extra attention during formulating any water management strategies for this region
3. Scenario analyses of probable future impacts of tile drainage in stream flow in the Red River basin of the North

Collection and processing data for the Red River of the North Basin on topography, land use, soils, meteorology, hydrology, and water quality have been completed. SWAT model was successfully set up and calibrated for the Western Wild Rice River watershed. A method for mapping the existing tile-drained areas in the upper Red River basin is being developed.

# **Co-entrapment of iron nanoparticles and trichloroethylene degrading bacteria in alginate biopolymer for groundwater remediation**

Project Number: 2011ND244B

Fellow: Sharanya Shanbhogue

Adviser: Achintya Bezbaruah and Eakalak Khan

Start Date: 3/1/2011

End Date: 2/29/2012

## **Publication:**

Bezbaruah, A.N.; Shanbhogue, S.S.; Simsek, S.; Khan, E. Encapsulation of Iron Nanoparticles in Alginate Biopolymer for Trichloroethylene Remediation, *Journal of Nanoparticle Research*, 13:6673-6681, 2011.

Shanbhogue, S.S.; Simsek, S.; Khan, E.; Bezbaruah, A. Encapsulation of Iron Nanoparticles and Trichloroethylene Degrading Bacteria for Groundwater Clean-up, *Proc. World Environmental and Water Resources Congress 2011*, pp. 1075-1082, 2011 (also oral presentation).

## **Progress:**

The focus of this research is on co-entrapment of microorganisms and NZVI to remediate contaminated groundwater. The combined effect of microorganisms and NZVI will increase the degradation efficiency and allow nearly complete degradation of target contaminants. The contaminant chosen to work on is trichloroethylene (TCE), a halogenated aliphatic organic compound and a universal degreasing agent. TCE is a Class A carcinogen and one of the most commonly detected volatile organic chemical in ground water. Bioremediation of TCE is one of the most attractive methods to remove TCE from the environment. The method of bioremediation has been used at an anaerobic aquifer at St. Joseph, Michigan, a TCE contaminated site on the National Priority List (NPL). The study of co-entrapment of NZVI and microorganisms is expected to yield critical conclusions for improving dechlorination reactions. An effective 2-stage remediation process will be engineered with the expectation that the contamination degradation efficiency will increase, reducing the contaminant to its very benign forms. The proposed new method has the potential to satisfy section 121 of CERCLA.

## **Project Objectives:**

The main objective of this study is the co-entrapment of NZVI and TCE degrading bacteria in alginate polymer to achieve complete degradation of TCE in groundwater. The Specific Objectives of this research are:

1. Encapsulation of NZVI and microorganism in alginate polymer.
2. Conduct degradation experiments for TCE
3. Monitor TCE degradation using different species of TCE degrading bacteria (e.g., *Pseudomonas putida* F1, *Dehalococcoides* sp.) in the capsule.

Alginate capsules have been synthesized and NZVI was successfully encapsulated. Diffusion and treatability studies using encapsulated NZVI was performed for varying concentrations of TCE. Microorganisms were encapsulated and used for TCE degradation. Encapsulated NZVI, microorganisms, and the combined metal-microorganism system were found to be effective for TCE degradation. The project is ongoing using different strains of microbes to test the efficacy of the combined metal-microorganism system. Based on the progress made so far a manuscript was submitted to a peer reviewed journal.

# **Role of agricultural drainage on transport of *Cryptosporidium* oocysts in North Dakota**

Project Number: 2011ND243B

Fellow: Tanush Wadhawan

Adviser: Eakalak Khan and John McEvoy

Start Date: 3/1/2011

End Date: 2/29/2012

## **Publication:**

Wadhawan, T., Kasi, M., McEvoy, J., Chu, M., Khan, E. 2012. Investigating transport of *Cryptosporidium* under snowmelt conditions. World Environmental and Water Resources Congress. Albuquerque, New Mexico, USA.

Wadhawan, T., Kasi, M., Chu, M., Khan, E., McEvoy, J. 2012. Bench-scale rainfall and snowfall simulations to understand *Cryptosporidium parvum* transport in subsurface groundwater regimes. IV International Giardia and *Cryptosporidium* Conference, Wellington, New Zealand.

Wadhawan, T., Kasi, M., McEvoy, J., Chu, M., Khan, E. 2011. Role of Manure Application on Soil in Preventing Groundwater Contamination by *Cryptosporidium*. 84rd Water Environment Federation's Annual Technical Exhibition and Conference, Los Angeles, USA.

Wadhawan, T., Khan, E., Chu, X., McEvoy, J. 2011. Adsorption of *Cryptosporidium parvum* oocyst on soil samples obtained from North Dakota. 2011 Land Grant & Sea Grant National Water Conference, Washington DC, USA.

## **Progress:**

The sources for *Cryptosporidium* can be human or animal; however, the understanding of a relationship between the source and disease transmission is limited. A study investigating the degree of strain variation exhibited by bovine and human isolates was conducted on the Red River of the North in 1998. Samples were collected from various parts of the watersheds for the presence of *Cryptosporidium* and identified 20 different isolates, some of which might cause human infections. *Cryptosporidium parvum* will be used as a model organism in this project for two reasons. First, the bovine strains of *C. parvum* were the most common isolates observed. Secondly, *C. parvum* is the primary cause of human cryptosporidiosis in the region. The main scope of this project is to investigate the role of agricultural drainage system on transport of *Cryptosporidium* in North Dakota. The specific objectives of this study are: 1) To investigate adsorption and desorption of *Cryptosporidium parvum* on the soils obtained from agricultural fields in North Dakota. 2) To determine the effects of agricultural drainage systems on the transport of *Cryptosporidium parvum* through the soils obtained from agricultural fields in North Dakota by simulating subsurface tile drains in a soil box and 3) To study the transport of *Cryptosporidium* found in the manure applied to a subsurface drained

agricultural field in North Dakota. This research is expected to demonstrate for the first time the role of agricultural drainage system on the transport of *Cryptosporidium*. It will greatly benefit North Dakota in identifying a possible source of cryptosporidiosis outbreaks.

To achieve objective 1, experiments were designed to investigate adsorption of *Cryptosporidium parvum* on natural soil by performing isotherm in the presence of manure. High adsorption of *Cryptosporidium parvum* was observed on both loam and clayey soil. Without manure, very slightly higher adsorption of *Cryptosporidium* to the clayey soil ( $1.94 \times 10^5$  to  $1.75 \times 10^6$ ) compared to the loam soil ( $1.92 \times 10^5$  to  $1.68 \times 10^6$ ) was observed. With manure, the adsorption of *Cryptosporidium* to the rocky sand increased by 35.2%-36.2%, while for the loam and clayey soil, the increases in adsorption were minimal, 2.3%- 2.6% and 1.3%-1.5%, respectively. The high adsorption of the oocysts to soil with or without addition of manure will prevent transport of *Cryptosporidium* and restrict oocysts to the top soil in the fields. The adsorption onto the rocky sand increased with the addition of manure. Addition of manure to the loam and clayey soil resulted in slight increase in the adsorption of the *Cryptosporidium*. Good fit of the experimental data to the Freundlich isotherm suggests the role of non-specific physical forces involved in the adsorption of oocysts. The Redlich-Peterson isotherm also gave a good fit to both of the soils. The Redlich-Peterson model fitted the experimental data as good as the Freundlich model. These isotherm results should be useful in drawing inferences for further studies on *Cryptosporidium* transport.

Objective 2 was achieved by performing soil box experiments. We evaluated the effect of two precipitation conditions with varying intensities and durations on *Cryptosporidium parvum* transport. The experiments were performed in a soil box with a surface area of 172 cm<sup>2</sup>. Chloride tracer or *C. parvum* was applied to the surface of a saturated soil box. Rainfall intensities of 2.5 and 5 cm/h for 1 h duration were simulated using a syringe pump. Conditions were chosen to represent 1 and 10 year rainfall events for Fargo, North Dakota, USA. To simulate snowmelt, the saturated soil box was kept at -20°C for 24 h, after which 200 g of ice (~116 mL of water) was applied. Two subsurface groundwater regimes drained the soil box using a drain tile or a small outlet. Both regimes were studied with and without the overland flow. *Cryptosporidium* was quantified using immunofluorescence microscopy. Our data show that none of the precipitation events could completely drain chloride (Figure 4) or *C. parvum* (data not shown). The numbers of *C. parvum* in the effluent varied from 5% to 50% of the influent. This study provides insight into the transport of *Cryptosporidium* under varying precipitation and subsurface flow conditions.

# **Temporal-spatial distribution (dynamics) of phytoplankton and diversity in relation to lake physical and chemical condition**

**Project Number:** 2011ND240B

**Fellow:** Veselina Valkov

**Adviser:** Wei Lin

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

## **Publications:**

Veselina Valkov "Temporal-spatial distribution (dynamics) of phytoplankton and diversity in relation to lake physical and chemical condition". Young Professionals **poster** competition at the North Dakota Water & Pollution Control Conference (NDWPCC), 11-13 October 2011, Bismarck

Veseline Valkov (2011) "Temporal-spatial distribution (dynamics) of phytoplankton and diversity in relation to lake physical and chemical condition". Presented at the 5th International Student Prairie Conference, June 2-3, 2011. North Dakota State University, Fargo, ND.

## **Progress:**

Eutrophication as a result of anthropogenic input of nutrients especially phosphorus and nitrogen, that cause real (or perceived) concerns for surface water quality, is significant problem for North Dakota lakes and reservoirs. The increased input of nutrients causes excessive and rapid growth of phytoplankton. Biological productivity is high and the diversity of biological populations may be limited. The massive development of phytoplankton especially Cyanophyceae (blue-green algae), tends to form dense "surface blooms". Some species are toxic or could cause allergic reaction. Surface blooms reduce light and nutrient availability to other algal species leading to lower algal diversity. High turbidity and floating films (mats) caused by algal blooms reduce light penetration, which is important for photosynthetic activity of submersed macrophytes. Sedimentation dead algae biomass causes increasing of organic matter in the sediments. Decomposition of sediment organic matter causes oxygen depletion, forming anoxic condition at the sediment surface and the deep water fauna becomes deprived of oxygen. Even pelagic fish, which release their eggs in open water to sink to the bottom, cannot continue to reproduce under anoxic sediment surface. Anoxic condition causes change in microbial and chemical processes. The release of nutrients from the anoxic sediments (internal loading) especially phosphorus causes self acceleration of eutrophication. Spectacular fish kill may result from advanced eutrophication. The goal of the proposed project is to expand existing research on water quality of the eutrophic impoundment Henrich Martin Dam. The scope is to understand how variation in physical and chemical disturbances influenced spatial - temporal phytoplankton distribution and diversity and can cause shifts in changes in community patterns.

Two sampling seasons: 6.4-10.15 2010 with aeration and 6.30-8.11 2011, with aeration/non-aeration period. Samples were taken from 5 sites on a biweekly basis. Physical parameters: Secchi depth, Turbidity, Temperature, DO, pH, Total and volatile suspended solids (TSS, VSS), Chemical oxygen demand (COD). Chemical parameters – NO<sub>2</sub>, NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub>, Total Nitrogen (TN), Total dissolved nitrogen (TDN), Total Phosphorus (TP), Total dissolved P (DP), Soluble Reactive Phosphorus (SRP) and Total reactive Phosphorus (TRP), Acid hydrolysable Phosphorus. Biological parameters/analyses - Chl a and phytoplankton abundance and speciation.

The water temperature increased steadily with maximum in July after that decreased temporally. The aeration system caused destratification, eliminating the thermocline. On July 13th, 2011, the aeration system was turned off and a week later a thermocline was developed between 1 and 2 m, with a drop of 3.20 °C. With depth, DO concentration decreases all seasons from super-saturated on the surface to close to standard of 5.00 mg/l on the bottom. In 2011, after aeration was stopped, a concentration of 1.75 mg/l was measured at just 2.00 m and dropped to 0.8 mg/l in the deepest layer. That could be correlated with algal accumulation on the surface, diurnal variation in respiration and photosynthesis, microbiological activity, wind mixing, however high organic matter in the water column. Diurnal fluctuation of DO 24-hour measurements without aeration shows highest values near the surface between 8-11 p.m. and minimum were observed in the morning's hours at 6 a.m. The dramatically depleted DO likely caused the observed fish kill, floating decayed biomass, scum and rotten smell on the shallow parts in the lake. The aeration system as a short-term affect maybe is not efficient to reach these outlying edges. High density macrophytes as *Contail* observed at the shallow parts also contribute to the DO balance. COD decreased in depth indicates high consumption of oxygen at the bottom of Site A during non-aerated period. The algae growth affected transparency of water. The high Secchi depth 1.70 -2.50 m. was contributed to low Chl a and TSS and VSS concentrations usually occurred in June 2010/2011. On June 18, 2010, the transparency decreased rapidly influenced from strong wind that increased TSS and turbidity and high Diatom dominance and Chl a reached 32.89 µg/l (Chl a 20-56 µg/l are classified as eutrophic). In July, 13th 2011, observed with necked eye healthy population of *Aphanizomenon* sp. (Cyanophyceae) aggregates decreased Secchi depth to 1.60 m. Two weeks after stopping aeration system high turbidity reduced the Secchi depth to 0.70 m. The *Aphanizomenon* aggregates were packed up and population constituted from single filaments. The algae cells that were brought out from upper layer from tabulation slowly migrated and accumulated on the surface and Chl a increased from 32.07 to 79.87 µg/l on A site, while in the bottom the change was less remarkable. Aeration system also affected nutrients variation in depth. During aeration period there were little concentration gradient in nutrients occur in water column. Inorganic nitrogen decreases in the summer remains low. In well oxidized waters, the release of NH<sub>4</sub> could be rapidly nitrified to NO<sub>3</sub> and further use from phytoplankton. The high values of NH<sub>4</sub> were detected late in summer as result of decomposition of prior high growth of algae, high temperatures and low DO. In addition, stopped aeration, very slow mixing, and oxygen depletion increased NH<sub>4</sub> in the bottom from 0.04 to 1.08 mg/l just after a month imply high rate of mineralization of organic matter and possible release from sediments. At



such low DO concentrations high organic matter in the sediments (about 13% measured) is oxidized with  $\text{NO}_3$  and is reduced to  $\text{N}_2$ . As a controlling algal growth factor SRP varies in concentration following the temporal variation in temperature that affecting oxidation-reduction of Al and Fe exchangeable complexes in the sediments. The high concentrations of SRP 0.05-0.18 mg/l were no limitation factor for phytoplankton growth. No clear correlation was found between the phytoplankton growth and SRP. In 2011 developed thermocline caused increasing concentration in the bottom layers up to 0.47mg/l, which also indicate sediment release as a main source of P in the HMD. The N limitation, decreased temperature, return to artificial mixing and seasonal turnover effect maybe led to reduced algae abundance, high transparency (Secchi disk = 3.6 m at A site and touched the bottom at C and D site). As result DO levels incased and the  $\text{NH}_4$  and SRP flux to the overlying surface water was sufficient for phytoplankton growth later in November. The strong N limitation in the summer months in all sampling seasons led to low the N:P ratio. The Redfield ratio 7.5:1 on biomass basis typically evaluated with respect to phytoplankton growth, which is empirically determinate values are needed for growth. Nuisance blue-green algae (Cyanobacteria) blooms are often associated with lakes that have low nitrogen to phosphorus (N:P) ratios.

During study periods, ten classes were identified, dominated of which were Bacillariophyceae, Dinophyceae, Chryptophyceae, Cyanophyceae and Chlorophyceae. The changing in the temperature, duration of the day and nutrient supply led to the succession sequence. At the beginning of June Bacillariophyceae was dominant class in terms of species numbers, density and biomass. Later in summer the Dinophyceae starts to be a dominating class. Class Chrysophyceae characterized with occurrence of mixotrophy was found once in June 2010 and in August and September in 2011, probably due to their requirements for high water temperature or nutrition supply. The period between these two dominant classes is characterized with more diversity between groups. Not as expected under Nitrogen limited condition class Cyanophyceae in 2010 contributed no more than 11% from the total abundance in summer, while in 2011 was 76%, in spite of aeration system still worked. The population of Cyanophyceae three weeks after aeration was stopped collapsed. Potential explanation include acceleration of buoyancy-induced vertical migration rates, protection against photo inhibition, reduce grazing losses, high oxygen, that could inhibit which could potentially inhibit nitrogenase activity, exposure to the nutrients and turbulence. The population continued to decrease and replaced by Dinoflagellate.

# Information Transfer Program Introduction

Information dissemination is done through an annual newsletter, and presentations and publications by grant and fellowship recipients. A web site also helps disseminating institute research information. The institute's website address is <http://www.ndsu.edu/wrri>. Past newsletters can be accessed through the institute web site. Technical reports of Fellowship projects authored by the Fellows and advisers are also placed on the institute web site.

## Information Dissemination and Communication

**Project Number:** 2011ND233B

**Start Date:** 3/1/2011

**End Date:** 2/29/2012

**Principal Investigator:** G. Padmanabhan

Activities to disseminate institute and other research under this project included:

1. Maintaining a web site
2. Publishing a newsletter
3. Publication of Fellowship and other research done through the Institute
4. Presentation of research results to state and federal water agencies
5. Sponsor or co-sponsor local or regional conferences
6. Sponsor special lectures on water-related topics of interest

The website of the Institute was updated at least quarterly, and more often when a research project wished to provide updates or when a Fellow graduates. The website provides additional details on the research. The list of Institute Affiliate Faculty with their expertise was updated. Research reports published by the institute were placed on this web site as and when they became available. The institute web site is <http://www.ndsu.edu/wrri>.

The Institute continued its annual newsletter, which highlights the graduate research fellowship program, the research grants associated with it, and general summaries of ongoing research. The newsletter profiled institute research and researchers and published other newsworthy water issues in the State. A copy of the newsletter can also be found on the institute web site: <http://www.ndsu.edu/wrri>

The Institute continued its efforts to enhance communication between the State and Federal agency personnel and university faculty and students. Advisors and fellows presented their research results to State and Federal professionals in Bismarck. The Institute also encouraged its Fellows and faculty to attend seminars and conferences held in the region. Modest support for travel was provided by the institute whenever appropriate.

## **NDWRRI sponsored the 2nd Annual Distinguished Water Seminar**

The 2<sup>nd</sup> Annual Distinguished Water Seminar was held on February 21, 2012 at NDSU campus open to faculty and students of NDSU and UND and the public. Dr. Kurt Fausch, a nationally and internationally known conservation biologist, conducted the seminar titled, " Linked for Life: the importance of sustaining hidden connections for conservation in streams." The talk was co-sponsored by the the Environmental and Conservation Sciences program, Agricultural and Biosystems Engineering department, Civil Engineering department, Soil Science department, and School of Natural Resources Sciences and Management, all of North Dakota State University. The seminar is the second of the annual Distinguished Water Seminar series by eminent water professionals on emerging issues, challenges and new research directions in water resources.

Dr. Fausch is a professor in the Department of Fish, Wildlife, and Conservation Biology at Colorado State University. He teaches Fish Conservation Biology and a graduate course in Community Ecology, and is active in the Graduate Degree Program in Ecology. His collaborative research has taken him worldwide, and especially to Hokkaido in northern Japan where he worked with colleagues over a 15-year period. These experiences were chronicled in the documentary film *RiverWebs*, directed and produced by Jeremy Monroe of *Freshwaters Illustrated*, which has been broadcast to more than 100 million homes nationwide on PBS. He has received several prestigious awards for his research and outreach, including the first International Fisheries Science Prize from the World Council of Fisheries Societies (2008) and Awards of Excellence from the American Fisheries Society (2010). He serves on the Independent Science Advisory Board of the Northwest Power and Conservation Council, which advises managers of the Columbia River about fish and wildlife conservation. Kurt is currently writing a book for a popular audience with the goal of engaging the public in understanding the interconnections between streams and rivers and their landscapes, and the importance of conserving these ecosystems.

Following the seminar, students and faculty had opportunity to interact with Dr. Fausch.

G. Padmanabhan, Director NDWRRI, Craig Stockwell, Director Environmental and Conservation Sciences Graduate Program, and Xuefeng Chu, Civil Engineering department were the organizing committee members.

**NDWRRI partnered with ND Department of Health, USGS, ND State Water Commission in organizing the ND Water Quality Monitoring Conference February 27-29, 2012 in Bismarck, North Dakota**

NDWRRI director, Dr. G. Padmanabhan, participated in the planning as a planning committee member. He encouraged several institute affiliate faculty and NDWRRI Graduate Fellows to participate in the planning and to present at the conference. Dr. Padmanabhan also moderated a session titled “Emerging Contaminants and Threats to Water Quality”. Seven past and current NDWRRI Fellows and eight institute affiliate faculty presented and participated at the conference.

Detailed agenda of the conference can be viewed at:

<http://info.bismarckstate.edu/ceti/waterquality/>

Presentation abstracts can be viewed at

<http://info.bismarckstate.edu/ceti/waterquality/pdfs/presentation-abstracts.pdf>

## **NATIONAL COMPETITIVE PROGRAM (104-G)**

Two proposals were submitted; but were not funded:

“Does the Longitudinal Profile and Distribution of Stream Power Affect the Efficiency and Ecology of Agricultural Drainage Ditches?” by Phil Gerla, Department of Geology and Geological Engineering, University of North Dakota

“Impacts of Land-use Change on Environmental and Water Quality in the James River Basin: Predicting Economic and Physical Outcomes for Efficient Policy Making” by Robert Hearne and Zhulu Lin, Agricultural Economics and Agribusiness, North Dakota State University

### **Institute Publications**

Effects of Major Flooding on Water and Sediment Characteristics in an Urban Environment, Adam Guy and Thomas DeSutter, Department of Soil Science, NDSU, NDWRRI report ND11-01, June 2011.

### **Publications by Institute Fellows and Faculty**

Leelaruban, N., P. Oduor, A. Akyuz, S. Shaik, and **G. Padmanabhan** (2012) Leveraging a Spatio-Temporal Drought Severity and Coverage Index with Crop Yield Modelled as a Stochastic Process, in International Journal of Hydrology Science and Technology (IJHST) Vol. 2 No. 3

**Kasi, M., McEvoy, J., Padmanabhan, G., and Khan, E.** (2011) Groundwater Remediation Using Enricher Reactor - Permeable Reactive Biobarrier for Periodically Absent Contaminants. Water Environment Research, 83(7), pp. 603-612.

Suman L. Shrestha, Xuelian Bai, David J. Smith, Heldur Hakk, **Francis X. M. Casey**, Gerald L. Larsen, and **G. Padmanabhan** (2011) Synthesis and Characterization of Radiolabeled 17 $\beta$ -estradiol Conjugates, Journal of Labelled Compounds and Radiopharmaceuticals, 2011, 54, 64. Also published online in Wiley Online Library, DOI: 10.1002/jlcr.1864

**Krajangpan, S.; Bezbaruah, A.N.;** Chisholm, B.J. Groundwater Arsenic Remediation using Amphiphilic Polysiloxane Graft Copolymer Coated Iron Nanoparticles, Proc. World Environmental and Water Resources Congress 2011, pp. 1083-1088, 2011 (also oral presentation).

**Shabnam, R.;** Simsek, S.; **Khan, E.; McEvoy, J.; Bezbaruah, A.** Diffusion and Treatability Studies with Biopolymer Encapsulated Zero-valent Iron Nanoparticles, Proc. World Environmental and Water Resources Congress 2011, pp. 1543-1551, 2011(also oral presentation).

**Kalita, H.**; Chisholm, B.; **Bezbaruah, A.N.** Novel Arsenic Ion-imprinted Polymer: Simultaneous Removal As(III) and As(V) from Water, Proc. World Environmental and Water Resources Congress 2011, pp. 3396-3401, 2011 (also oral presentation).

**Simsek, H.**, **Kasi, M.**, **Wadhawan, T.**, Bye, C., Blonigen, M., and **Khan, E.** (2011) Modeling a Two-Stage Trickling Filter Wastewater Treatment Plant to Simulate the Fate of Dissolved Organic Nitrogen and Its Biodegradability. In Proceeding 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA, Water Environment Federation.

**Kasi, M.**, **Simsek, H.**, Blonigen, M., and **Khan, E.** (2011) Fate of Bioavailable and Biodegradable Dissolved Organic Nitrogen in a Two-Stage Trickling Filter Wastewater Treatment Plant. In Proceeding 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA, Water Environment Federation.

**Wadhawan, T.**, **Kasi, M.**, **McEvoy, J.**, **Chu, X.**, and **Khan, E.** (2011) Role of Manure Application on Soil in Preventing Groundwater Contamination by Cryptosporidium. In Proceeding 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA, Water Environment Federation.

#### **Conference/Seminar Presentations**

**Kasi, M.**, **Simsek, H.**, **Wadhawan, T.**, Bye, C., Blonigen, M., and **Khan, E.** (2011) Modeling a Two-Stage Trickling Filter Wastewater Treatment Plant to Simulate the Fate of Dissolved Organic Nitrogen and Its Biodegradability. Presented at the 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA. (Presented by M. Kasi)

**Kasi, M.**, **Simsek, H.**, Blonigen, M., and **Khan, E.** (2011) Fate of Bioavailable and Biodegradable Dissolved Organic Nitrogen in a Two-Stage Trickling Filter Wastewater Treatment Plant. Presented at the 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA. (Presented by H. Simsek)

**Wadhawan, T.**, **Kasi, M.**, **McEvoy, J.**, **Chu, X.**, and **Khan, E.** (2011) Role of Manure Application on Soil in Preventing Groundwater Contamination by Cryptosporidium. Presented at the 84th Annual Water Environment Federation Technical Exposition and Conference, Oct. 15-19, 2011, Los Angeles, CA. (Presented by T. Wadhawan)

**Krajangpan, S.**; **Bezbaruah, A.N.**; Chisholm, B.J. Polymer Coated Zero-valent Iron Nanoparticles for Arsenic Reduction, The NDWPCC Strom Water Conference, Bismark, ND, October 2011.

**Krajangpan, S.; Bezbaruah, A.N.**; Chisholm, B.J. Groundwater Contaminant Removal using Nanosized Zero-valent Iron Coated by Amphiphilic Polysiloxane Graft Copolymer, The 5th Annual Joint Student Environmental Conference, Fargo, ND, June, 2011.

**Patent**

**Bezbaruah, A.**; Chisholm, B.; **Krajangpan, S.** Polymeric Delivery Vehicle for Nanoparticles. Nationalized PCT Patent, 2011/0042,325, Feb 24, 2011.

***THESES AND DISSERTATIONS***

Qigang Chang (Ph. D)  
Ishara Rijal (M. S)

Abstracts of their theses follow:

## ABSTRACT

Chang, Qigang, Ph.D., Environmental and Conservation Sciences Program, College of Graduate and Interdisciplinary Studies, North Dakota State University, August 2011.  
Iron-Impregnated Granular Activated Carbon for Arsenic Removal from Drinking Water.  
Major Professor: Dr. Wei Lin.

Iron-impregnated granular activated carbon (Fe-GAC) can effectively remove arsenic from water. A new multi-step iron impregnation method was developed in this study to impregnate GAC with high amount of iron that possesses desired characteristics: stable, even distribution, and high arsenic adsorption capacity. Research was carried out to investigate the impact of the amount of impregnated iron on arsenic adsorption properties: capacity, affinity, and kinetics.

Fe-GACs with different iron contents were characterized in terms of the amount, stability, distribution, morphology, and species of impregnated iron. It was found that high amount of iron was stably impregnated in GAC. Scanning electron microscopy (SEM) and energy dispersive X-ray spectroscopy (EDS) analysis demonstrated that the impregnated iron was evenly distributed on the internal surface of GAC. Impregnated iron formed nano-size particles, and existed in both crystalline (akaganeite) and amorphous iron forms.

Arsenic adsorption tests were conducted using Fe-GACs with iron content of 1.64-28.90% and adsorption isotherms covered the range of arsenic equilibrium concentration that is typical for drinking water treatment. Langmuir model fit both As(III) and As(V) isotherm adsorption well. The amount of impregnated iron affects arsenic maximum adsorption capacity ( $q_m$ ), but have little impact on the Langmuir constant  $b$  (the affinity of adsorbent for adsorbate). The  $q_m$  for both As(V) and As(III) adsorptions increased significantly with increase of the amount of impregnated iron up to 13.59%. Further increase of iron amounts caused gradually decrease of  $q_m$  for As(V). The  $q_m$  for As (III) continued to slightly increase to 2456  $\mu\text{g/g}$  at iron content of 24.73%. BET analysis indicated that the surface area of Fe-GAC decreased with increase of the iron amount. At 13.59%, impregnated iron fully covered the internal surface area of GAC.

A new second-order kinetic model was developed to investigate the impact of the amounts of impregnated iron on arsenic adsorption kinetics. This new kinetic model can well fit arsenic adsorption kinetics. With iron content increase from 1.64% to 28.90%, the intrinsic adsorption rate constants kept reducing from  $4.6 \times 10^{-2}$  1/hr to  $1.18 \times 10^{-3}$  1/hr, which indicates that impregnated iron slows arsenic intraparticle diffusion rate in Fe-GAC. Decreased arsenic intraparticle diffusion rate was most likely caused by reduced pore size of Fe-GACs. The intrinsic adsorption rate constant is not dependent on initial concentration in kinetic tests, which eases the comparison of the arsenic adsorption kinetics of Fe-GACs.

Groundwater sample taken from a former superfund site in North Dakota with a total arsenic concentration of 205  $\mu\text{g/L}$  was used in column tests to investigate the effect of the amount of impregnated iron on arsenic breakthroughs. With empty bed contact time



(EBCT) of 600 second, the number of bed volume (BV) treated at breakthrough ( $10 \mu\text{g/L}$ ) increased from 140 to 1000 with iron content increasing 4.56-13.59%. Further increase of iron content, BV treated at breakthrough slightly decreased. The slope of breakthrough curves became smaller with increase of iron content, which implies that arsenic intraparticle diffusion rate in Fe-GACs decreased with more impregnated iron. Fe-GACs with high iron content require long EBCT to achieve a better performance. This indicates that Fe-GACs with high iron content require longer EBCT to utilize Fe-GACs in column tests.

## ABSTRACT

Ishara Rijal, M.S., Agricultural and Biosystems Engineering, College of Engineering and Architecture, North Dakota State University, August 2011. Reference Evaporation and Actual Evaporation Measurements in Southeastern North Dakota. Major Professor: Dr. Jia Xinhua.

Subsurface drainage (SSD) has been used to remove excess water from field in the upper Midwest US for more than a century, but only since last decade in the Red River Basin of North in North Dakota. The water leaving from a SSD system can affect both the quality and quantity of water which flows to a surface water system. Therefore, determination of the water balance components is the first step to study the impact of SSD on water quantity, while evapotranspiration (ET), one of the most important components in the water balance, needs to be accurately measured for SSD field. A field experiment was conducted to study the water balance in SSD and undrained (having no artificial drainage system) fields in southeast ND. The field had three different water management systems: 22 ha undrained (UD), 11 ha subsurface drained and the remaining 11 ha subsurface drained and subsurface irrigated. The ET rates were measured directly using an eddy covariance (EC) system for the SSD and UD fields. The changes in water table were monitored in 8 wells installed in both fields. Rainfall, SSD drainage volume, and soil moisture at six different depths at two locations were measured in both fields. The measurements were conducted in the growing seasons of 2009 and 2010. The ET rates were calculated for two different field crops: Corn (*Zea Mays*) in 2009 and soybean (*Glycine Max*) in 2010. Crop coefficient ( $K_c$ ) value was also developed using the ET measured by the EC system and the reference ET ( $ET_{ref}$ ) estimated using the American Society of Civil Engineers Environmental and Water Resources Institute (ASCE-EWRI, alfalfa) method. The  $ET_{ref}$  was also estimated using the ASCE-EWRI grass and the Jensen Haise (JH) methods.