


# Mendel, the Chromosomal Theory of Inheritance, and DNA As the Hereditary Material

## Why Do We Sequence Nucleic Acids??

- Dumb question???
  - Yes, because we all know that nucleic acids control phenotype.
    - Well we have not known that forever.
  - So how did we learn this?

## Three lines of evidence converged that lead to this discovery

- Phenotypes are controlled by genes
- Genes are located on chromosomes
- Chromosomes are made up of DNA (one of the nucleic acids).



## Gregor Mendel

### Dramatically changed our perception of heredity

- Particulate factor concept
  - *Some physical factor existed that controlled phenotype*

## Traits have a dominant and recessive forms

- Proof
  - F<sub>1</sub> generation
    - Dominant form appears
    - Recessive form disappears
- F<sub>2</sub>
  - Recessive form reappears

## Mendel's 1st Law, the Law of Segregation

- A single form of the factor controlling phenotype was passed to the gamete during reproduction.
  - Event occurs during reduction step of meiosis
- One of two forms of the factor was passed through the gamete to the offspring.
  - Proof??
    - F<sub>2</sub>
      - 3:1 ratio in F<sub>2</sub> generation segregating for one trait
        - 3/4 dominant form
        - 1/4 recessive form
    - F<sub>3</sub> generation
      - Offspring of recessive F<sub>2</sub> plants all recessive form
      - Some offspring of F<sub>2</sub> dominant form plants all dominant form
      - Some offspring of F<sub>2</sub> dominant form plants produce 3:1 dominant to recessive forms
      - Ratio of dominant form F<sub>2</sub> plants in F<sub>3</sub> generation
        - 2/3 segregate for dominant and recessive forms in 3:1 ratio
        - 1/3 all dominant F<sub>3</sub> plants

## **Mendel's 2nd Law, the Law of Independent Assortment**

- Each trait was controlled by a unique factor
- Proof??
  - 9:3:3:1 ratio in the F2 generation segregating for two traits
    - The cross product of two 3:1 ratios is 9:3:3:1

## **Mendel**

- DID not consider the actual physical entity that controls experiments
  - Others discovered that entity

## **Other experiments determined**

- Mendel's factors (genes) reside on chromosomes
- DNA was the heredity material.

## **Naming the Mendelian Laws**

- **Correns (1900)**
  - Referred to segregation and assortment
- **Morgan (1916)**
  - First to use the terms:
    - ***Law of Segregation***
    - ***Law of Independent Assortment***

## Genes Reside on Chromosomes



### Eduard Strasburger (1876)

- Cell division is a universal activity of all higher organisms
  - Same process is observed in plants




### Walther Flemming (1878)

- Structures had a string like appearance to them
  - Termed the structures chromatin (or colored substance)
- Also developed the concept of cell division
  - ***Called cell division mitosis.***




### **Edouard van Beneden (1883)**

- Egg cell + sperm cell fertilization
  - Resulting cell has the diploid chromosome number
- Sperm and egg cell
  - Each contribute equal numbers of chromosomes to the zygote
- Discovered meiosis



### **Heinrich Wilhelm Gottfried von Waldeyer-Hartz (1888)**

- Called the structures dividing during mitosis chromosomes
  - **Chromosomes (=colored bodies)**



### **August Weismann (1892)**

- Germline Theory
  - Sperm and egg cells
    - Contain exactly half the number of chromosomes
      - ***Transmit hereditary information***
  - Somatic cells
    - Carry out normal body functions



Hugo deVries (1889 on)

### Performed hybridization experiments and discovered

- ***Each trait controlled by a different factor***
  - Observed 3:1 ratio in F<sub>2</sub>
- Called the hereditary factor a ***pangene***
- Individual pangenes controlled all traits
  - Pangenes located in the cell of a diploid organism

### Initially unaware that Mendel proved this earlier

- Later tried to publish without referencing Mendel
  - Correns corrected him
    - ***Admitted that Mendel was the first to discover the laws of genetics***

### Considered pangenes

- Larger than a single chemical molecule
  - But still invisible

### Reproductive cells

- Receive half of the pangenes during meiosis
  - When reproductive cells unite
    - Diploid number of pangenes is restored

***First linkage between inheritance and reproductive cells***



### Carl Correns (1900)

- Study inheritance in plants
  - Published results in paper
    - “G. Mendel's Law Concerning the Behavior of the Progeny of Racial Hybrids”



### Erich Tschermak (1900)

- Plant breeder working on wheat, barley, and oats
  - Tried to combine earliness and high yield
    - Considered the "father" of Austrian plant breeding
- Did genetic experiments with pea
  - **Referenced Mendel in his publication**

### Rediscovering Mendel's Concepts of Genetics

- Referenced in publications by:
  - DeVries (April 1900)
  - Correns (May 1900)
  - Tschermak (August ??? 1900)



## Chromosome Theory of Inheritance (1902-1903)

*Chromosomes are the carrier of Mendelian factors and meiosis is the basis of separating the factors into gametes.*



### Theodor Boveri (1902)

- Observed
  - All male and female chromosome must be present to develop a functioning organism
  - Linked chromosomes and the factors that were described by Mendel
- Quote
  - *"... the characters dealt with in Mendelian experiments are truly connected to specific chromosomes."*



### Walter Sutton (1902)

- Described chromosomes as unique individual units
  - That occur in pairs
  - Separate during meiosis
- Quote
  - *Chromosomes "...may constitute the physical basis of the Mendelian law of heredity."*

# Linking Genes and Chromosome

## Thomas Hunt Morgan and Calvin Bridges



### Thomas Hunt Morgan (1910)

- Discovered a mutant white eye *Drosophila*
  - Different than the wild type red eye
- Performed genetic experiments
  - ***Results proved the eye color gene was located on the X chromosome***



### Calvin Bridges (1914)

- Studied Morgan's white eye mutant
  - Coupled the presence of the X chromosome with a specific eye color
  - ***Conclusively demonstrated genes indeed reside on chromosomes***

### **Problem with this concept!!**

- Chromosomes carried the genetic information
  - They must contain all the genetic factors
- But, the number of chromosomes is less than the number of traits.
  - ***Now it was essential to show chromosomes contain many factors***

### **Solution: Multiple Genes Reside on Chromosome!!**

- **Sturtevant, Bridges, Morgan (1919)**
  - Mated among *Drosophila* with several different contrasting phenotypes
    - Multiple genes are organized into a linear linkage group
      - ***Number of linkage groups equals the number of chromosomes***

**It could now be stated:**

***All features necessary for a hereditary unit are found in chromosomes!!!***

## Position Effect



### Sturtevant (1925)


- If the physical environment of a gene is altered
  - Expression of the gene is affected
- Therefore
  - *Physical structure of the chromosome is essential for the correct phenotypic expression*

### From a modern perspective

- This is the first solid evidence that we should take a **genomics approach** to fully understand gene expression.
- Therefore
  - *Completely characterizing (=sequencing) all of the genetic material in the cell is necessary.*

## SO WHAT IS THE GENETIC MATERIAL???

# History of DNA As the Genetic Material



**Ernst Haeckel (1866)**

- Nucleus transmitted hereditary information to the next generation



**Friedrich Miescher (1871-1874)**

## **Studied pus cells collected from bandages from surgeries**

- Collected white blood cells
- White blood cells primarily composed of nuclei
  - *Called this nuclear material **nuclein***

## **Determined that nuclein contained two classes of chemicals**

- **Acidic component**
  - *Now we know the component is **DNA and RNA***
- **Basic component**
  - *Now know that is **histone proteins***

## Linking DNA and Heredity




### Fred Griffith (1928)

- Worked with lethal and non-lethal strains of the *Streptococcus pneumoniae*
  - Converted a non-lethal strain to a lethal strain
- Conversion involved
  - Mixing dead lethal and live non-lethal strains

### Griffith's Transforming principle

- Converts one phenotype to another
  - This is the true nature of a gene
  - **These are two alleles of the same gene**
- **So what is the chemical nature of the transforming principle???**



Oswald Avery   Colin MacLeod   Maclyn McCarty

### Avery, MacLeod, and McCarty (1944)

- Transforming principle
  - DNA was the transforming principle (from the acidic component)
  - Not protein or RNA
    - The other two constituents in the nucleus.

### RNA Is Also A Genetic Material



- Heinz Fraenkel-Conrat (1957)
  - RNA viruses exist
    - Interconverted strains of tobacco mosaic virus
      - RNA mediated the interconversion and can be a genetic material

# Chemical Structure of DNA



## James Watson and Francis Crick (1953)


- DNA is double-stranded
- Strands are oriented in an anti-parallel manner to each other
- Purines nucleotides are opposite pyrimidines nucleotides
  - Guanine hydrogen bonds with cytosine
  - Adenine hydrogen bonds with thymine
- Structure is stabilized by
  - Hydrogen bonds
  - Hydrophobic bonding between stacked bases

## Watson and Crick

- Did not perform any experiments
  - Based on research of others



## Research results of others that aided Watson and Crick



- **Erwin Chargaff**
  - Concentrations of guanine and cytosine were always equal in DNA
  - Concentrations of adenine and thymine were equal in DNA



- **Rosalind Franklin and Maurice Wilkins:**
  - Used X-ray crystallography to study structure of DNA

## Watson and Crick major contributions to describing the structure of

- DNA had a repeating structures (nucleotides)
- DNA was of a constant width
- DNA was double-stranded