

POST EXPOSURE PROPHYLAXIS FOR NONOCCUPATIONAL EXPOSURE TO HIV (HIV PEP)

Pharmacy Assessment Protocol

ELIGIBILITY FOR SERVICE

Patients who meet the following criteria are eligible for post-exposure prophylaxis:

- Potential exposure is non-occupational in nature
- Potential exposure occurred after a single high-risk event within the **past 72 hours**
 - Exposed to bodily fluids from a person with HIV/reasonably suspected to have HIV
- Age 13 years and older
- Weight >77 lbs
- Not pregnant/breastfeeding
- Are not currently taking HIV PrEP therapy
- Without known current acute/chronic kidney disease and/or decreased kidney function

The pharmacy will refer those not meeting the above criteria for further medical evaluation.

TESTING CRITERIA

- Anyone eligible for HIV PEP should be tested with a CLIA-Waived HIV Test
- If a patient refuses the test and meets eligibility criteria, prescribing can still occur

PRESCRIBING

Upon a negative HIV rapid test, the pharmacist shall identify any precautions or contraindications to therapy and then prescribe the most appropriate treatment:

TREATMENT			
MEDICATION	Frequency	Duration	Notes
emtricitabine 200 mg/ tenofovir disoproxil fumarate 300mg (Truvada® or generic)	Once daily	28 Days	Dosing adjustments required with renal dysfunction if CrCL <60 ml/min.
PLUS			
raltegravir 400mg	Twice daily	28 Days	
	OR		
Dolutegravir 50mg	Once daily	28 Days	

FOLLOW-UP

If the patient tests positive for HIV upon provision of a CLIA-Waived test, the patient will be immediately referred to further medical evaluation. Patients prescribed PEP should be referred for follow up with a primary care provider to obtain the following as soon as possible:

Serum creatinine, ALT/AST, Hepatitis B screening, Hepatitis C screening, pregnancy test (if applicable)

The patient should be contacted within 4 to 6 weeks to obtain a follow-up HIV screening test.